THE OPIOID SAFETY INITIATIVE AND VETERAN SUICIDES

Joshua Tibbitts and Benjamin W. Cowan*

February 24, 2021

Abstract: We investigate the relationship between opioid diverting policy and suicides among the veteran population. The opioid epidemic of the past two decades has had devastating health consequences among U.S. veterans and military personnel. In 2013, the Veterans Health Administration (VA) implemented the Opioid Safety Initiative (OSI) with the goal of discouraging prescription opioids dispensed to VA patients. Between 2012 and 2017, prescription opioids dispensed by the VA fell 41% (VA, 2018). Because this involved the aggressive curtailing of opioid prescriptions for many VA patients, OSI may have had a detrimental effect on veterans' mental health leading to suicide in extreme cases. In addition, because rural veterans have much higher rates of VA enrollment, prescription opioid use and related abuse, and lower rates of substance abuse and mental health treatment utilization, we expect any effect of OSI on veteran suicides to be concentrated in rural areas. We find that OSI raised the veteran suicide rate relative to the civilian rate with rural veterans suffering the lion's share of the increase. In particular, OSI raised the rural veteran suicide rate by roughly one-third within the first year of implementation (2013) and by 45 percent by 2016.

JEL Codes: I12; I18; D11; D12

Keywords: Opioids, Policy, Suicides, Veteran

* Joshua Tibbitts is a doctoral candidate at Washington State University and Benjamin Cowan is an Associate Professor in the School of Economic Sciences at Washington State University and a Research Associate at the National Bureau of Economic Research. We thank Jill McCluskey, Shanthi Manian, and seminar participants at Washington State University and the Health Education Action Lab at Southern Utah University for helpful comments. Any errors are our own. Correspondence: Joshua.Tibbitts@wsu.edu and ben.cowan@wsu.edu.

1. Introduction:

Over the last few decades, the United States has seen a surge of substance abuse disorders related to opioids. From 1999 to 2018, almost 450,000 people died from an opioid-related drug overdose; of the 67,367 fatal drug overdoses in 2018, 70 percent were opioid-related (WONDER, 2020). The broad detrimental effects of this crisis have been heterogeneous. For example, individuals living in rural areas are 87 percent more likely to receive an opioid prescription compared with individuals in urban areas (García, 2019); this corresponds with elevated opioid-related poisonings rates, which for rural residents have increased at 3 times the rate of those who live in metropolitan areas (Keyes, et al., 2014). Veterans have also been particularly susceptible to this crisis with rural veterans faring worse than their urban counterparts (Johnson et al., 2015; McCarthy et al., 2012; Mohamed et al., 2009; Wallace et al., 2006).

In response to the opioid epidemic, state policymakers have implemented Prescription Drug Monitoring Programs (PDMPs) to limit the supply of prescription opioids by deterring patients from "doctor shopping" and enhancing physician accountability in their opioid prescribing practices. The implementation of PDMPs has resulted in successful supply reductions of prescription opioids in recent years (Bao et al., 2016; Deyo et al., 2018; Haffajee et al., 2018; Kilby, 2015; Moyo et al., 2017; Suffoletto et al., 2018). However, there is also evidence that these policies led to an increase in illicit opioid abuse and fatalities (Alpert et al., 2017; Delcher et al., 2016; Kilby, 2015; Meinhofer, 2017)

The Veteran Affairs Health System (VA)--which includes roughly half of the total veteran population, including 93 percent of service-connected veterans (i.e., those injured during their service) (NCVAS, 2020), as well as many current military personnel--did not utilize state PDMP databases or create their own national prescription history database until 2013. The number of individuals treated with opioids at the VA increased at a staggering rate from the turn of the century until it hit a peak in 2012 (see Figure 1); of all veterans receiving opioid prescriptions, more than one-third were for longer

than 90 days ("chronic use") (VA, 2015). The high percentage of long-term opioid prescriptions among veterans is associated with a high incidence of opioid substance abuse disorder and opioid-related mortality (Baser et al., 2014; Bohnert et al., 2011; Axelrod, 2013).

In an attempt to combat the growing opioid crisis among veterans, the VA implemented the Opioid Safety Initiative (OSI), which was primarily designed to curtail opioids prescriptions, especially high dose prescriptions (a daily dose of more than 100 morphine milligram equivalents, or MME's), long-term prescriptions, and concomitant prescriptions (e.g. opioids and benzodiazepines) for comorbid maladies (Sandbrink, 2019; GAO, 2018; Good, 2017; VA, 2020). Curtailment initiatives included establishing a database of patients' prescription histories (with a "clinical leader" at each facility tracking database metrics to promote safer prescribing; see Lin et al., 2017), increasing drug screening among patients, and encouraging physicians to substitute toward alternative pain management practices through VA instituted education programs and directives. Initially launched in five small regions in 2012[1] and nationally implemented in 2013, this policy has added additional directives[2] in successive years, including in 2016 when the VA began mandating queries into state PDMP databases for their physicians (prior to that, queries into state PDMP databases were encouraged for the physician, but they were not mandatory). Coinciding with the rollout of the policy, the number of opioids dispensed immediately began to diminish. From 2012 to 2017, the VA's opioid prescription rate, or the number of unique patients prescribed an opioid, decreased by 41 percent (VA, 2018); as of the third quarter in the year 2020, the decrease was 64 percent compared to 2012 (VA, 2021), indicating that the VA continues to aggressively limit the use of prescription opioids among veterans.

[1] These regions were located in Minnesota, which is a state that is not included in our dataset.
[2] See Figure 2, which shows a timeline of OSI implementation and additional directives.

In this paper, we examine the effects of OSI on veteran and military personnel suicides. Despite the policy's apparent success in reducing the number of opioids dispensed, there have been accusations from within and without the VA Health System[3] that OSI policy encouraged physicians to "cut off" veterans from their pain medications (GAO, 2018). The curtailment of opioid prescriptions may increase the risk of suicide due to mental and physical anguish related to withdrawal and physical pain (Borgschulte et al., 2019; Demidenko et al., 2017; Kuramoto et al., 2012). Because of the high incidence of chronic pain with corresponding high rates of opioid prescriptions, the veteran population is acutely susceptible to these increased suicide risks (Oquendo and Volkow, 2018; Johnson et al., 2015). Thus, it is possible that in an attempt to aid the veteran population by curtailing opioids, OSI unintentionally increased suicides among veterans and military personnel.

We investigate this question with a triple difference (DDD) empirical model. Our outcome is county-level suicide rates per 100,000 individuals in both civilian and veteran/military personnel populations. OSI only affected VA users, so civilians serve as a control group in our analysis. We do not have information on which veterans and military personnel receive healthcare through the VA, so all veterans are included in our treatment group and the effects we estimate are likely smaller than what we would find for VA patients specifically.[4] We further differentiate civilian and veteran populations by urban-rural status. The urban-rural distinction is made due to the vast differences in health care options, treatment availability, opioid prescription rates, and cultural norms between these veteran populations; these dissimilarities should result in differential impacts of OSI on suicide rates among rural and urban veterans, as we argue in Section 1.3. We exploit the national implementation

[3] In 2011, prior to national OSI implementation, the Minneapolis, St. Cloud, and Fargo VAs pioneered OSI, resulting in reduced high-dose opioid prescriptions by 90% prior to 2014; long-term opioid prescriptions were reduced by 78%. The Star Tribune, a Minneapolis newspaper published "Cut Off – Veterans Struggle to Live with VA's New Painkiller Policy," which quotes several addiction specialists accusing the VA of creating addicts and then cutting off their supply. Additionally, the Government Accountability Office (GAO) alleged that the VA failed to establish "safe" tapering programs: https://www.gao.gov/assets/700/692061.pdf.

[4] Veterans and current military personnel who are referred to the VA by their health provider, TRICARE, along with their dependents, are the only population affected by this policy; the vast majority of VA users are veterans.

of OSI in 2013 to identify the effect of OSI on rural and urban veterans relative to their civilian counterparts.

Other studies point to the discontinuance of prescription drugs as a risk factor for suicide, but these studies do not establish a causal link between the two (Demidenko et al., 2017; Kuramoto et al., 2012; Oliva et al., 2020). As we describe in the next section, veterans and military personnel are a population particularly susceptible to increased suicide risk resulting from a loss of prescription analgesics (Toblin et al., 2014; Johannes et al., 2010; Cesur et al., 2019; Baser et al., 2014; Kang et al., 2015).

We first show that prior to the implementation of OSI, trends in suicides are similar between veterans/military and civilians. However, after the implementation of OSI, the suicide rate for veteran/military personnel increases relative to the civilian rate, and this effect is concentrated among rural veterans. In the year of implementation, rural veteran suicides increased roughly one-third relative to the pre-treatment mean; in 2016, the total increase in the rural veteran suicide rate was 45 percent. Urban veterans also experienced statistically significant increases in suicides in 2015 and 2016 on the order of 15 percent. These findings are robust to controlling for different sets of time-variant explanatory variables and modeling choices, as we show in Section 3.

1.1 Background: Veterans and Opioids

Perhaps no other group has suffered the ill effects of opioid substance abuse as United States veterans and military personnel. Opioid-related overdose mortality rates for veterans are 1.3 to 2.1 times higher than for civilians (Bohnert et al., 2011; Axelrod, 2013) and the prevalence of substance abuse disorders for veterans is almost seven times higher (Baser et al., 2014). In 2016, it was estimated that approximately 65,000 veterans were treated for opioid addictions (VA Opioid Prescription Policy, 2015) and the overall overdose rate per 100,000 veterans had risen 50 percent compared to 2010 (Lewei, et al., 2019).

Due to their unique training and experiences (stressful simulations and combat operations), physical and psychological pain is purported to be a primary contributor to the opioid crisis within veteran communities (Bennett et al., 2013; Cesur et al., 2019). An estimated 45% of veterans suffer from chronic pain (Clancy, 2015; Sandbrink, 2017), a rate that is 3 to 5 times higher compared to the civilian population (Toblin et al., 2014; Johannes et al., 2010). Among veterans who receive treatment at VA facilities, over 50% suffer from chronic pain (Gironda et al., 2006). Because of these higher pain levels, as of 2011, nearly half of the veteran population diagnosed with non-cancer related pain were prescribed an opioid; 40 percent of those prescriptions were for long term use (Edlund et al. 2014), which increases the likelihood of addiction relative to those with limited opioid exposure (Edlund et al., 2010).

Veterans who have been involved in combat operations--approximately 2.1 million since 2001 (Wenger et al., 2018)--have a particularly high incidence of physical and psychological pain, which is a strong antecedent to opioid-use disorders (Cesur et al., 2019). Chronic pain is the most common medical issue for veterans returning from combat operations, and nearly 60% of returning veterans suffer from it (VHA, 2013). In the aftermath of major combat operations in Iraq and Afghanistan, the rate of opioid use disorders among veterans has increased by 55% (VA Opioid Prescription Policy, 2015). Combat-related injuries and psychologically traumatic experiences (e.g. witnessing death) play a particularly important role in the rate of opioid use disorders among veterans (Cesur et al., 2019). Operation Iraqi Freedom (OIF) and Operation Enduring Freedom (OEF) has also exposed millions of servicemembers to cheap access to opium during their military deployments; this phenomenon, coupled with the VA's historically lax opioid prescription policies, has significantly contributed to opioid dependence and abuse among the population of veterans and military personnel (Cesur et al., 2019).

Because many veterans suffer from comorbid physical and mental maladies (physical trauma and Post Traumatic Stress Disorder, or PTSD), especially those veterans who have combat experience, concurrent use of prescription opioids and Central Nervous System (CNS) depressants are relatively common; 27% of veterans who were prescribed opioids were also jointly prescribed benzodiazepines, a CNS depressant (Park et al. 2015). The joint use of opioids and benzodiazepines increase the probability of emergency room visits and inpatient admission due to opioid overdose (Sun et al., 2017). The presence of both types of medications has been present in roughly half of all veteran opioid overdose deaths (Park et al. 2015).

1.2 Background: The Opioid Safety Initiative (OSI)

To mitigate the destruction caused by overprescribing opiate-derived analgesics to millions of veterans, the Department of Veteran Affairs (VA), acknowledging their role in fueling this opioid epidemic among veterans, instituted the Opioid Safety Initiative (OSI) nationwide in 2013. This was followed by "Academic Detailing" initiatives in 2014, which provided opioid prescribing education for physicians affiliated with the VA.[5] Next came the Comprehensive Addiction and Recovery Act (CARA) in 2016, which mandated that all physicians within the VA conduct annual inquiries into state PDMP databases at least once a year for their patients (although physicians were encouraged to query state PDMP databases from the start of OSI, it was not mandated) (GAO, 2018; Good, 2017).

The mechanisms by which the VA has reduced prescribed opioids includes national opioid databases ("OSI Dashboards"), which utilize the VA's Electronic Health Records and includes state PDMP databases. Physicians are able to track their patients' medication history, urine drug screening, and alternative forms of chronic pain therapy (Petzel, 2014). The VA tracks "key clinical indicators," which include the number of unique patients dispensed an opioid, the number of patients on

[5] Good (2017) finds that physicians who engaged in Academic Detailing decreased their prescribing of high-dose opioids by 58 percent compared with a 34 percent reduction among physicians who did not.

concomitant prescriptions of benzodiazepines and opioids, unique patients prescribed long-term opioid therapy, and mean daily opioid dosages. Using these metrics, high-risk patients are identified (Petzel, 2014).

Since the implementation of OSI, VA physicians were strongly encouraged, through newly instituted pain management guidelines, to use alternative means to treat pain and to avoid opioid prescriptions where possible (VA, 2014). Physicians within the VA who do not apply the prescribing guidelines initiated by OSI are identified and may be subject to VA counseling, education, or other actions taken by the VA (Petzel, 2014). Thus, physicians may feel pressure to improve VA-tracked opioid metrics to avoid internal investigations and actions. There are multiple reports of physicians within the VA who have rapidly tapered their patients off of opioids--a potentially risky outcome (Demidenko et al., 2017; Dubin et al., 2017)--and who have set arbitrary dose limits for stable patients who were on chronic opioid treatments (Gellad et al., 2017; Good, 2017). Notably, as published in the VA/Department of Defense (DOD) Clinical Practice Guideline for Management of Opioid Therapy for Chronic Pain, physicians were instructed to taper opioid therapy patients at a rate of 20-50 percent per week (VA, 2010), which is a much higher rate than the 10 percent per month suggested by the CDC (CDC, 2016).

OSI has been highly successful in reducing opioid prescriptions within the VA. Since OSI was chartered by the Under Secretary for Health in August of 2012 after being piloted by several regions in Minnesota (Clancy, 2015), opioid-related prescriptions have steadily decreased in each successive fiscal quarter, including high dose prescriptions, long-term prescriptions, concomitant prescriptions of opioids and benzodiazepines, and the number of unique patients dispensed an opioid (Sandbrink, 2019). Although OSI was not fully implemented nationally until August of 2013, physicians likely anticipated the forthcoming changes starting in August of 2012 and appear to have

begun changing their opioid prescribing practices sometime in the fourth quarter of 2012 (See Figure 1).

1.3 Background: Veterans and Suicides

Over the same time period as the U.S. opioid epidemic grew, suicides also increased at a troubling rate. Prior to the year 1999, suicide rates experienced consistent annual declines in each of the previous 13 years (Curtin, et al., 2016). The age-adjusted suicide rate in the year 2017 was 33% higher when compared to the rate in 1999; between these years, the suicide rate increased in each successive year (Hedegaard, 2018). Suicides involving opioids, i.e. intentional fatal opioid poisonings, were twice as high in the year 2014 as they were in 1999 (Braden et al., 2017). Individuals who suffer from Opioid Use Disorder (OUD) are 13 times more likely to die by suicide (Wilcox et al., 2004) and prescription opioid misuse is significantly associated with suicidal ideation, suicide planning, and suicide attempts (Ashrafioun et al., 2017). Thus, opioid use and abuse may have a causal link to suicides. Opioid-related mortality and suicides are suggested to be two primary causes of the rising mortality rates among middle-aged white non-Hispanic men and women in the United States--a trend reversal after decades of progress (Case and Deaton, 2015).

A common factor in opioid use and suicide is physical pain. As stated earlier, physical pain initiated the opioid epidemic through the wide prescribing and misuse of opiate-derived analgesics before the detrimental effects were widely known (Van Zee, 2009; CBHSQ, 2017). Physical pain is also highly correlated with suicide ideation, suicide attempts, and suicide completion (Fishbain, 1999; Fishbain, 1991; Hooley, 2014; Tang and Crane, 2006), though suicide risk does not appear to be dependent on a specific injury or type of chronic pain (Ilgen et al., 2013). Pain may elevate the risk of suicide both indirectly through prescription opioid use and directly through one's diminished quality of life. Although opioids can potentially enhance one's quality of life by alleviating pain, when

individuals' use of opioids become abusive, prescription opioid use appear to increase the risk of suicide, on average.

As in the opioid epidemic, the population of veterans and military personnel are acutely vulnerable to the current suicide crisis. Compared to the general population, veterans' risk of suicide is 41 to 61 percent higher (Kang et al., 2015). Because veterans have extremely high rates of opioid substance abuse and mental and physical pain (including PTSD for those veterans deployed in combat operations), veterans are highly susceptible to the risk of suicide (Bryan et al., 2015; Maguen et al., 2011; Kim et al., 2012).

Although opioid substance abuse is associated with an increased risk of suicide, forced opioid discontinuation may exacerbate this risk, especially within the first three months of discontinuation (Oliva et al., 2020; Talari and Yara, 2020); evidence from VA patients during the year of OSI implementation suggests that the risk of suicide may not stabilize until 6 to 12 months following opioid discontinuation (Oliva et al., 2020)[6]. Citing evidence from an "alarming increase in reports of patient suffering and suicide" due to forced opioid tapering within the VA, the International Stakeholder Community of Pain Experts and Leaders recommend that policies aimed at prohibiting or minimizing forced opioid tapering be enacted, along with other recommendations (Darnall et al., 2019). OSI encourages physicians to reduce opioid prescriptions and implement tapering programs (VA, 2010); those veterans with mental health and substance abuse disorders are more likely to endure opioid tapering and discontinuation within the VA (Oliva et al., 2020; Vanderlip et al., 2014). An estimated 75 percent of all clinician-initiated opioid discontinuation within the VA were the result of substance

[6] The authors also show that the risk of suicide following opioid discontinuation is exacerbated with the length of time engaged in opioid treatment. The hazard ratios for patients who committed suicide after stopping opioid treatment ranged from 2.02 for patients on opioids less than 30 days to 7.99 for patients on opioids 400 days or more (Oliva et al., 2020).

abuse-related aberrant behavior exhibited by the patient (Lovejoy et al., 2017)[7]. Additionally, there are concerns that physicians have initiated forced opioid tapers to get patients under a dosage threshold for the primary purpose of improving OSI performance measures (Gellad et al., 2017).

1.4 Background: Rural-Urban Differences in Potential Impacts of OSI

According to the VA's Office of Rural Health (ORH), approximately 4.7 million veterans live in rural communities in the United States, which represents 30 percent of the total veteran population (NOSORH, 2014). In comparison, rural civilians only account for 19 percent of the total civilian population (VA's ORH, 2020). Not only do rural veterans make up a comparatively large portion of the total veteran population, but they also have high VA utilization rates. Nearly 60 percent of rural veterans receive their health care from the VA (NOSORH, 2014); of this population, nearly 55 percent suffer from at least one service-connected condition.[8] The rate of VA utilization among rural veterans is much higher than the 37 percent enrollment rate for urban veterans (VA's ORH, 2020). This is one reason we expect VA policy to disproportionately affect rural veterans compared with urban veterans.

Not only do rural veterans have a higher percentage of VA-enrolled users, but this population has also been more adversely affected by the concurrent opioid and suicide crises. When compared to their urban counterparts, veterans living in rural communities are prescribed over 30 percent more opioids per capita; most of these differences between urban-rural opioid prescriptions were for long term use, which OSI was focused on curtailing (VA, 2011). After the implementation of OSI, total rural opioid prescriptions fell by 36 percent (Lund et al., 2019). In addition to having higher per capita rates of opioid prescriptions (Hudson et al., 2017; Lund et al., 2019), rural veterans are also more likely

[7] 85 percent of all veterans on long-term opioid therapy were forced to discontinue their opioid regimen (Lovejoy et al., 2017).

[8] The rate of VA enrollment among service-connected disabled veterans is over 93 percent (VA, 2017) https://www.va.gov/vetdata/docs/Quickfacts/VA_Utilization_Profile_2017.pdf. With a high rate of VA enrollment and correspondingly high rates of chronic pain and opioid prescriptions, this population may be heavily impacted by OSI.

to suffer fatal opioid overdoses (GAO, 2018). This is possibly because rural veterans receive medication assisted treatment and other evidenced-based treatments for addiction at a lower rate than urban veterans, despite their higher rates of substance abuse (Edmonds et al., 2020; GAO, 2018; Smalley et al., 2010).

In terms of suicide, the rural veteran population is at higher risk (by 20 - 22 percent) compared to urban veterans, even after controlling for availability of mental health services, such as distance to mental health treatment centers (McCarthy et al., 2012). Numerous studies have shown that high rates of suicide and comparatively low rates of treatment for rural veterans[9] have less to do with treatment availability and are more likely determined by economic circumstances, comparatively lower quality of life, and rural culture (McCarthy et al., 2012; Mohamed et al., 2009 Teich et al., 2017; Wallace et al., 2006). Oliva et al. (2020) found that rural veterans are at a greater risk of suicide following opioid discontinuation than urban veterans. In sum, the rural veteran population has higher VA utilization rates, opioid prescription rates (which have dramatically decreased following the implementation of OSI), overdose rates, rates of suicide, and lower addiction and mental health treatment utilization rates, compared with their urban counterparts. For these reasons, we allow for different estimates of the effects of OSI on suicides by urban-rural status in our regression analysis.

2. Data

For our main dependent variable of interest, suicides, we obtained data from the Center for Disease Control's National Violent Death Reporting System's Restricted Access Database (CDC-NVDRS-RAD) which links data from vital records, coroner/medical examiners, and law enforcement agencies in participating states. This dataset is the most comprehensive data available for violent deaths

[9] Many studies show that rural veterans are less likely than their urban counterparts to utilize mental health services, which includes services for substance abuse disorders.

including: suicides, homicides, unintentional firearm-related deaths, and violence-related deaths where the intent cannot be determined (NVDRS, 2018). The restricted-access database includes veteran status and county of residence for all reported deaths. 17 states began participating in NVDRS prior to OSI implementation in 2013, and these are the ones we include in our sample (other states' data became available only after OSI was implemented and are left out). We have data on 16 states starting in 2010 and data on Ohio starting in 2011. When we leave Ohio out of our analysis to have a perfectly balanced panel, our results are very similar to those shown in the paper (available upon request). For this paper, we aggregated the data to the county-year level.

The NVDRS codes a death as a suicide if intent can be determined. Because this dataset only includes "violent" deaths, accidental overdoses or overdoses in which intent cannot be determined are not accounted for; thus, suicides may be undercounted to the extent that some overdoses are purposeful but intent cannot be determined. If a county is not represented in the database for a specific year, it is because there is no record of a suicide occurring in that county-year. With this in mind, we add a zero for all counties in participating states that were not represented in the NVDRS-RAD database in a specific year.

We gather population, demographic, and socio-economic variables for each county-year for both veterans and non-veterans from the American Community Survey (ACS). These variables are discussed in the next section. Prior literature on opioid supply shocks and suicide (Borgschulte, et al., 2019) shows that policies aimed at reducing the supply of opioids, when combined with geographical proximity to addiction treatment centers, act to reduce suicide. Therefore, following Borgschulte et al. (2019) and Swensen (2015), we used NAICS codes 621420 and 623220 from the Consumer Business Patterns (CBP) to identify inpatient and outpatient treatment centers in each county-year.

We use the rural-urban continuum code developed by the US Department of Agriculture Economic Research Service (USDA, 2013) to determine the rural-urban distinction for counties in

our analysis. Counties are classified based on 9 categories that are determined by their population size and whether they are adjacent to a metropolitan county. As in Kaplan et al. (2012), we assigned counties as urban for classification codes 1-3 (metropolitan counties) and rural for codes 4-9 (non-metropolitan counties).

Finally, to control for other prescription opioid policy supply shocks over our sample period, we gathered user access date data for Prescription Drug Monitoring Programs from Borgschulte, et al., (2019), the Prescription Drug Abuse Policy System (PDAPS), and the National Alliance for Model State Drug Laws (NAMSDL). For must-access PDMP information, we use data published by Borgschulte, et al., (2019), Buchmueller and Carey (2018), and the New Jersey Prescription Monitoring Program (NJPMP).

3. Empirical Strategy

Due to the large share of veterans and military personnel who suffer from comorbid ailments (i.e. chronic pain and PTSD) leading to substance abuse disorder, our hypothesis is that veteran opioid users' suicide choices will be affected by the negative supply shock to prescription opioids due to OSI. For reasons described above, we expect that rural veterans will be more affected by OSI than urban veterans. To investigate these possibilities, we use a generalized triple difference (DDD) regression model. Our treatment group is veterans and military personnel, and our control group is non-veterans/civilians. The treatment policy is OSI, which only affects those who receive healthcare through the VA. The vast majority of these are veterans/military personnel, though a tiny fraction of civilians receive care through the VA, such as immediate family members (VA, 2017). Some veterans do not receive healthcare through the VA, but we do not observe source of healthcare in our suicide data, and the choice to receive care through the VA may be endogenous with respect to OSI. Thus, we estimate the effect of OSI on the incidence of suicide among all veterans/military personnel

relative to non-veterans/civilians (we refer to the former group simply as "veterans" and the latter group as "civilians" below). We then allow the effect of OSI to also vary by whether a suicide victim lived in a rural county.

The specific regression equation in our analysis is as follows:

$$\left(\frac{suicide}{100K}\right)_{vct} = \theta_c + \theta_t + \delta_1 * veteran_v + \delta_2 * veteran_v * rural_c + \sum_{t=2011}^{2016}(\alpha_t * veteran_v) + \sum_{t=2011}^{2016}(\gamma_t * rural_c) + \sum_{t=2011}^{2016}(\rho_t * veteran_v * rural_c) + X_{vct}\pi + \varepsilon_{vct}. \quad (1)$$

The unit of observation in our study is a county by year by veteran status. θ_c and θ_t represent county and year fixed effects, respectively; county fixed effects control for time-invariant differences by county, and year fixed effects control for secular changes underlying the determinants of suicide. The coefficients δ_1 and $\delta_1 + \delta_2$ measure baseline (2010) differences in the suicide rate of urban and rural veterans, respectively. The α_t terms measure how the suicide rate evolves for urban veterans, relative to urban civilians, over time (2011-2016). The γ_t terms measure how the suicide rate evolves for rural civilians relative to urban civilians over time. Finally, the ρ_t term measures how the suicide rate evolves for rural veterans relative urban veterans, all relative to the same urban-rural difference in civilian suicide rates. The treatment year is 2013, which is the year that OSI was nationally implemented; we are interested in seeing how the parameters α_t, γ_t, and ρ_t evolve prior to and after 2013. This specification allows us to view the annual evolution of OSI's effect on veteran suicides, by rural status, where we are interested in the comparative evolution before and after OSI implementation; this is a generalization of the classic triple difference estimator (DDD).

X_{vct} represents a vector of time variant controls including an indicator for whether a county is located in a state with a PDMP and a separate one for a mandatory PDMP, the number of

treatment centers (sum of outpatient and inpatient) in a county, interactions between PDMP indicators and the number of treatment centers; and ACS estimates for both veteran and civilian populations within the county for: percent of the population aged 18-34 (the age group for which the risk of suicide is greatest), percent female, percent black, the unemployment rate, median income, and percent that have some college experience. We also include a full set of interactions between veteran status as well as rural county status and all other controls in the model.[10] All regressions are weighted by the total county population and standard errors are clustered by county.

In addition to a linear specification, like Borschulte et al. (2018), we estimate our model using a Poisson Quasi-Maximum Likelihood Estimator (QMLE), where the dependent variable is the count of suicides for each county and year. Moksony and Hegedűs (2014) argue that a Poisson regression is the most appropriate specification when estimating the variation of rare events such as suicides (e.g., a large number of "0" observations). Because our data is over-dispersed, we use the Poisson QMLE, which relaxes the assumption of mean-variance equality. With this specification, we offset veteran/military and civilian suicides by their respective populations; thus, our dependent variable is the log of the suicide count.

4. Trends and Descriptive Statistics:

Graphical analysis in Figure 3 based on raw suicide rates shows that although the suicide rate is roughly twice as high for veterans compared to civilians, trends in suicide rates for civilians and veterans are similar prior to 2013. This suggests that civilians offer a reasonable counterfactual for veterans and military personnel in a DD framework. Trends for these groups begin to diverge in 2013, the year of

[10] Veteran median income was missing for 380 counties in 2015. These missing values were replaced by the average veteran median income in 2014 and 2016 in their respective counties.

national OSI implementation; the percentage change in suicides per 100,000 individuals from 2012 (the year before OSI) to 2016 is 42 percent for veterans and 23 percent for civilians.

Table 1 provides pre-treatment descriptive statistics for veterans and non-veterans by rural/urban status. The differences in suicides per 100,000 between veterans and civilians shown in this table are consistent with the literature (Castro and Kintzle, 2014; Kang et al., 2015; Sher and Yehuda, 2015). The vast majority of veterans are male, and as a group they are much older than civilians. Bachelor's completion rates are similar across groups, and veterans fare somewhat better on average in terms of labor-market indicators (the unemployment rate and median income), likely owing somewhat to their age and gender profile.[11]

When looking at veterans and civilians across urban-rural status, we see that rural veterans/military personnel and civilians have worse outcomes in terms of suicide, economic well-being, and addiction treatment availability (at least in terms of the number of treatment facilities) when compared to their urban counterparts. For example, the mean suicide rate per 100,000, median income, and unemployment for rural veterans is 42.68, $30,860, and 8.14 percent, respectively; whereas urban veterans suffer from 33.88 suicides per 100,000, earn $39,318 in median income, and have a 7.18 percent unemployment rate. These differences are consistent with the literature cited above.

Table 2 provides descriptive statistics for both veterans and civilians differentiated between periods prior to and post-OSI implementation. The veteran suicide rate per 100,000 prior to OSI was 32.7 and increased to 43.44 after implementation (a 33 percent increase), whereas the civilian suicide rate increased from 16.65 prior to OSI to 18.52 after OSI (an 11 percent increase). Additionally, the large increase in the pre-post OSI suicide rate among veterans does not at first blush appear to be a

[11] Poorer economic conditions are associated with higher suicide rates (Denney et al., 2009; Tondo et al., 2006), but at least based on raw averages this would not seem to be the dominant explanation for the veteran-civilian suicide gap.

product of declining economic circumstances, as there is little change in average income and unemployment that occurred from before to after OSI implementation. This suggests that there may be other factors that are driving the relatively stark increase in veteran/military suicides.

5. Results

5.1 Baseline Results

Our main regression results are displayed in Table 3, where we present the comparative impact of OSI on suicides among veterans/military personnel and civilians. Column 1 in this table shows OLS estimates without any additional controls (X_{vct}) included in the regression equation and column 2 shows the coefficient estimates with the additional controls described in Section 3. Estimates for each of our four groups (urban/rural, civilian/veteran) are plotted for the specification with controls in Figure 4.

OLS estimates without controls show no consistent trend in urban civilian (base group) suicides until after 2013, when suicides increase modestly relative to 2010. When controls are included (column 2), these yearly differences shrink and become statistically insignificant. Rural civilians (see the coefficients on year*rural) experience no statistically significant yearly differences relative to urban civilians when controls are included in the model. When controls are not included, coefficient signs switch from negative to positive and a few are individually significant at the 10% level, but the coefficients remain relatively small and there is no clear trend in magnitudes after 2010.

With respect to the differential evolution for urban veteran suicides relative to urban civilian suicides, estimates with and without controls tell a very similar story: veteran suicides are elevated in all post-2010 years, but in 2015 and 2016 the effects jump in magnitude substantially and become statistically significant at the 5% level or better. In the case of column 2 (controls included), the effect in 2015 is 3.39 suicides per 100,000 (an 11% increase relative to the pre-treatment mean) and the effect

in 2016 is 3.93 suicides (13%). We note that we do not see immediate effects of OSI (implemented in 2013) when it comes to urban veteran suicides. Because of lower VA enrollment and opioid prescription rates among this group, it may be that effects of OSI become evident only after the policy has had a chance to grow in scope, which it did from 2013 to 2016 as opioid-related prescribing continually decreased coinciding with Academic Detailing and the implementation of CARA.

Our triple difference estimates, which measure any difference in suicides that rural veterans experience with respect to OSI compared to urban veterans, all relative to the same difference among civilians, are displayed at the end of Table 3. In this case, the findings are quite stark: while year effects prior to 2013 are small and statistically insignificant, they jump greatly in magnitude in 2013 and remain similar through 2016. In 2013, the difference in suicides between rural and urban veterans (relative to the urban-rural civilian difference) increased by 11.1 per 100,000 compared with 2010. This represents an increase of 32% relative to the pre-treatment mean for rural veterans. Effects in subsequent years are similar (albeit somewhat smaller) in both specifications. It is important to note that these effects are on top of estimated effects for all veterans, which increase themselves in 2015 and 2016. This is evident in Figure 4, which shows that rural veteran suicides continue an upward trajectory after 2013.

The immediate, large effect of OSI on rural veteran suicides in 2013 occurs even though the policy was officially implemented only in August of that year. We believe the reason for this is twofold: first, OSI was chartered by the Under Secretary for Health in August of the previous year after being piloted by several regions in Minnesota (Clancy, 2015). Therefore, physicians were aware that OSI was to be implemented nationwide in 2013. Additionally, from Figure 2, we see that unique patients dispensed an opioid reached a maximum point in the fourth quarter of 2012. Starting in the first quarter of 2013, we begin to see reductions in VA opioid prescribing that pre-dated the national implementation of OSI. Hence, we have reason to believe that VA physicians were aware of the policy and likely began changing their opioid prescribing behavior in anticipation of OSI throughout

the first 7 months of 2013. Second, as described above, suicide risks following opioid discontinuation are highest within the first three months (Demidenko et al., 2017). Furthermore, veterans who were initially forced to taper or discontinue their opioid prescriptions were often the most vulnerable: those with unstable addictions or who had displayed substance abuse aberrant behavior (Lovejoy et al., 2017). These factors combined suggest the immediate impact of the policy we observe for rural veterans is plausible.

The growing effects of OSI on both urban and rural veteran suicide rates over our sample period may be due to the evolution of OSI policy over time. After implementation of OSI in 2013, VA prescription opioid rates, including unique patients receiving opioids, high-dose prescription opioids, concomitant prescribing of opioids and benzodiazepines, and long-term prescription opioids, decline in each successive year; the drastic decrease in these opioid-related metrics, which the VA monitored after OSI implementation, coincide with a greater emphasis on Academic Detailing, which educates VA physicians on VA norms for opioid-related prescribing. As opioid prescriptions continued to decrease, including by way of more tapering and discontinuation programs, an increasing number of veterans likely became affected by the policy, which resulted in a greater number of veteran suicides.

5.2 Poisson Results

Table 4 shows the estimation results from the Poisson QMLE specification both without additional controls (column 1) and with additional controls (column 2) as discussed in Section 3. In this specification, urban civilians (base group) do not see an increase in suicides prior to 2014. Starting in 2014, the suicide trend for urban civilians is upward throughout the remainder of our sample, which is similar to what we find in our baseline OLS specification with no additional controls. When it comes to rural civilians, column 1 shows little different relative to urban civilians but column 2 (with

additional controls) suggests that rural civilian suicides are declining relative to urban civilian ones over the course of our sample period.

The main difference between our OLS and Poisson QMLE results is for urban veterans. Whereas we see distinct increases in suicides for urban veterans (relative to civilians) in 2015 and 2016 with our OLS specification, we see no such effects in the Poisson results. However, the triple difference estimates for the Poisson QMLE, which measure any difference in suicides that rural veterans experience compared to urban veterans, all relative to the same difference among civilians, are quite similar to their OLS counterparts. Starting in 2013 and continuing through 2016, these effects are large and statistically significant at the 5% level or better, ranging from about 30-44 percent depending on the year and whether additional controls are included. One caveat is that differences between 2012 and 2013 are smaller and not statistically significant at conventional levels in the Poisson QMLE, whereas they are in our baseline OLS results.

5.3 Threats to Identification

A potential threat to our identification strategy is the possibility that veteran migration across urban/rural areas coincides with or is caused by OSI. For example, rural veterans may seek additional healthcare services (including substance abuse treatment) to deal with the loss of opioids by migrating to urban areas. Or, as OSI is an opioid diverting policy, it is feasible that rural veterans may migrate to urban communities to access a more robust illicit drug market as a substitute for lost prescription opioids. If such migration occurred according to underlying factors that affect suicide risk, it would call into question our interpretation of results.

To investigate this phenomenon, we regress the percentage of each county's population who are veterans on time and year fixed effects and the interaction between year and rural indicators; the regression was weighted by county veteran population and the standard errors were clustered by county (see Table 4). The results indicate that although veteran populations are declining over time

in general (likely due to their much higher age than the civilian population, on average), this decline is no larger in rural areas, as all year by rural interaction terms are essentially zero. This suggests that differential migration across the urban-rural divide is not driving our results.

5.4 Discussion

In this sub-section, we attempt to place our results in the literature on opioid policy and suicides. Borgschulte et al., (2018) found that state PDMP policies have not increased suicides among the white non-Hispanic population over 30 years old. Furthermore, the authors found that when PDMP policy was paired with greater treatment center availability, suicides actually decreased. In contrast to the results of Borgschulte et al., (2018), we find that opioid diverting policy, specifically OSI, has increased suicides among the rural veteran population substantially and perhaps the urban veteran population more modestly.

We posit that the differences in our results with those in Borgschulte et al., (2018) are due both to the different populations analyzed as well as the mechanisms of each policy (PDMP's versus OSI). First, as mentioned above, United States veterans are perhaps the population that is most adversely affected by the opioid epidemic. This is due to their higher rates of chronic pain, greater access to prescription opioids, higher rates of substance abuse disorder, lower treatment utilization, and higher rates of suicide; comparing these adverse consequences between urban and rural veterans, rural veterans fare worse still. Second, OSI went beyond creating a database for patients' prescription histories or even mandating that physicians access that database (as was typically the case with PDMP's); it introduced additional measures that led to rapid opioid tapering and discontinuation programs that have been shown to increase the risk of suicide among a vulnerable population.

Another question in the literature is to what extent marijuana legalization policies provide a safer alternative (marijuana) to opioids when it comes to pain management. Anderson et al. (2014) showed that the suicide rate among males aged 20-29 (30-39) decreased by 9.2 to 10.8 (9.4 to 13.7)

percent in states that legalized medical marijuana compared with states that did not. The authors posit that these estimates are biased towards zero due to measurement error and are therefore a conservative estimate.

Lastly, we aim to put our results in context by approximating the ratio of suicides caused by OSI to opioid prescriptions foregone as a result of OSI. This involves a back-of-the-envelope calculation based on our results as well as several figures taken from other papers in the literature. At the peak of VA opioid prescribing in 2012, the number of unique veterans dispensed a prescription opioid was 1,017,826. By 2016, that number had fallen to 838,888, representing a decrease of 213,938 (Hadlandsmythe et al., 2018). Because an estimated 2.7 million rural veterans are enrolled in the VA while 7.9 million urban veterans are, and because rural veterans are prescribed 30 percent more opioids per capita, we estimate that 33 percent of the decrease in opioid patients was composed of rural veterans. Furthermore, our own results indicate that rural veteran suicides increased an average of 11.69 per 100,000 from the years 2013 to 2016 as a result of OSI (this figure is obtained from averaging the coefficients on the veteran by year effects and the rural by veterans by year effects for 2013-2016). Because there are approximately 4.7 million rural veterans overall, average annual suicides among the rural veteran population as a result of OSI is a little over 549 suicides, or 2,198 suicides as a result of OSI from 2013 to 2016. Therefore, for every 100 patients who were not prescribed opioids as a result of OSI, we estimate that there were 3 suicides. Following the same methodology for urban veterans (average increase in suicides per 100,000 following OSI of 2.78, representing a total of 1,621 urban veteran suicides from 2013-2016), the ratio of suicides to opioid patients foregone is approximately 0.01.

6. Conclusions

In this study, we investigate the link between VA supply-reducing opioid policy (OSI) and suicide among veterans and current military personnel. We find evidence that OSI resulted in an increase in veteran suicides. These effects appear to have grown through 2016 (the last year for which we have data) coinciding with additional OSI initiatives following initial national implementation in 2013 and a continual decline in opioid prescription at the VA. Our results show that rural veterans specifically have been particularly affected by OSI, with suicides increasing 45 percent relative to the pre-treatment mean by 2016. These differences are consistent with higher rates of VA enrollment, opioid use, and suicide risk among rural veterans as we have discussed.

The opioid crisis has had tremendous negative consequences among veterans; lower-bound estimates for annual health care costs for veterans with combat exposure alone are upwards of $1.04 billion for prescription painkillers and $470 million for heroin use (Cesur et al., 2019); the total opioid-related economic burden for the general population in the United States in 2013, which includes all veterans, was an estimated $78.5 billion (Florence et al., 2016). OSI, which reduced opioid prescribing for both those who entered the VA under the looser former opioid regime as well as newer veterans, may have great long-term benefits if it prevents opioid use disorders from occurring in the first place. However, our results suggest there are large benefits of effective countermeasures among rural veterans in the shadow of OSI. The most successful interventions would reduce veteran suicides while continually exposing fewer veterans to potentially dangerous pain mitigation strategies via opioid prescriptions.

Although it is beyond the scope of this paper, we surmise that successful treatment interventions may occur through enhanced addiction treatment networks. The rural veteran population may especially benefit from this, as they face significant barriers to treatment, which results in lower rates of treatment utilization compared to their urban counterparts (Cully et al., 2010; Edmonds et al., 2020; GAO, 2018; McCarthy et al., 2012; Mohamed et al., 2009; Possemato et al.,

2018; Teich et al., 2017). In 2016, the VA's Office of Connected Care, which focuses on improving rural veteran health care through technology, began distributing video-enabled computer tablets to veterans who suffer from mental health disorders and faced barriers to access mental health care due to their place of residence. Using the tablets to conduct telehealth appointments, veterans who were given these tablets had higher psychotherapy utilization rates and managed their medications more effectively compared to veterans who did not receive them (VA, 2018). Borgschulte et al. (2018) find that with a sufficient number of treatment centers in a county, suicides decline after the implementation of PDMP's. These results hint at strategies to prevent veteran suicides in the wake of opioid-diverting policy.

References

Alpert, Abby & David Powell & Rosalie Liccardo Pacula, 2017. "Supply-Side Drug Policy in the Presence of Substitutes: Evidence from the Introduction of Abuse-Deterrent Opioids," NBER Working Papers 23031, National Bureau of Economic Research, Inc.

Anderson, D. M., Rees, D. I., & Sabia, J. J. (2014). Medical marijuana laws and suicides by gender and age. *American journal of public health*, *104*(12), 2369–2376. https://doi.org/10.2105/AJPH.2013.301612

Ashrafioun, L., T. M. Bishop, K. R. Conner, and W. R. Pigeon. 2017. Frequency of prescription opioid misuse and suicidal ideation, planning, and attempts. Journal of Psychiatric Research 92:1–7.

Axelrod, Jim. 2013. "Veterans Dying from Overmedication," CBS News. Available at: https://www.cbsnews.com/news/veterans-dying-from-overmedication/

Bao, Y., Pan, Y., Taylor, A., Radakrishnan, S., Luo, F., Pincus, H. A., & Schackman, B. R. (2016). Prescription Drug Monitoring Programs Are Associated With Sustained Reductions In Opioid Prescribing By Physicians. *Health affairs (Project Hope)*, *35*(6), 1045–1051. https://doi.org/10.1377/hlthaff.2015.1673

Baser, Onur, Lin Xie, Jack Mardekian, David Schaaf, Li Wang, and Ashish V. Joshi. 2014. "Prevalence of Diagnosed Opioid Abuse and its Economic Burden in the Veterans Health Administration," Pain Practice 14(5): 437-445.

Bennett, A. S., Elliott, L., & Golub, A. (2013). Opioid and other substance misuse, overdose risk, and the potential for prevention among a sample of OEF/OIF veterans in New York City. *Substance use & misuse*, *48*(10), 894–907. https://doi.org/10.3109/10826084.2013.796991

Bohnert, Amy SB, Mark A. Ilgen, Sandro Galea, John F. McCarthy, and Frederic C. Blow. 2011. "Accidental Poisoning Mortality Among Patients in the Department of Veterans Affairs Health System," Medical Care 393-396.

Bohnert, K. M., Ilgen, M. A., Louzon, S., McCarthy, J. F., and Katz, I. R. (2017) Substance use disorders and the risk of suicide mortality among men and women in the US Veterans Health Administration. *Addiction*, 112: 1193–1201. doi: 10.1111/add.13774.

Borgschulte, M., Corredor-Waldron, A., and Marshall, G. (2018) A path out: Prescription drug abuse, treatment, and suicide. *Journal of Economic Behavior & Organization*, Volume 149, Pages 169-184, ISSN 0167-2681, https://doi.org/10.1016/j.jebo.2018.03.006.

Braden JB, Edlund MJ, Sullivan MD. Suicide Deaths With Opioid Poisoning in the United States: 1999-2014. *Am J Public Health*. 2017;107(3):421-426. doi:10.2105/AJPH.2016.303591

Bryan, C. J., Griffith, J. E., Pace, B. T., Hinkson, K. , Bryan, A. O., Clemans, T. A. and Imel, Z. E. (2015), Combat Exposure and Risk for Suicidal Thoughts and Behaviors Among Military Personnel and Veterans: A Systematic Review and Meta-Analysis. Suicide Life Threat Behav, 45: 633-649. doi:10.1111/sltb.12163

Buchmueller, Thomas C., and Colleen Carey. 2018. "The Effect of Prescription Drug Monitoring Programs on Opioid Utilization in Medicare." *American Economic Journal: Economic Policy*, 10 (1): 77-112.DOI: 10.1257/pol.20160094

Case, A. and Angus Deaton Rising midlife morbidity and mortality, US whites, Proceedings of the National Academy of Sciences Dec 2015, 112 (49) 15078-15083; DOI: 10.1073/pnas.1518393112

Castro CA, Kintzle S. Suicides in the military: the post-modern combat veteran and the Hemingway effect. Current Psychiatry Reports. 2014 Aug;16(8):460. DOI: 10.1007/s11920-014-0460-1.

Center for Behavioral Health Statistics and Quality (CBHSQ). *2017 National Survey on Drug Use and Health: Detailed Tables*. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2018.

Clancy C. Statement of Dr. Carolyn Clancy M.D. Interim Under Secretary For Health Veterans Health Administration (VHA) Department of Veterans Affairs (VA) Before the Committee on Veterans' Affairs United States Senate. 2015 March; 1-10. https://www.veterans.senate.gov/imo/media/doc/VA%20Clancy%20Testimony%203.26.20151.pdf

Cully, J.A., Jameson, J.P., Phillips, L.L., Kunik, M.E. and Fortney, J.C. (2010), Use of Psychotherapy by Rural and Urban Veterans. The Journal of Rural Health, 26: 225-233. doi:10.1111/j.1748-0361.2010.00294.x

Curtin SC, Warner M, Hedegaard H. Increase in suicide in the United States, 1999–2014. NCHS data brief, no 241. Hyattsville, MD: National Center for Health Statistics. 2016.

Darke, S. and Ross, J. (2002), Suicide among heroin users: rates, risk factors and methods. Addiction, 97: 1383-1394. doi:10.1046/j.1360-0443.2002.00214.x

Darnall, Beth B., David Juurlink, Robert D Kerns, Sean Mackey, Brent Van Dorsten, Keith Humphreys, Julio A Gonzalez-Sotomayor, Andrea Furlan, MD, Adam J Gordon, Debra B Gordon, Diane E Hoffman, Joel Katz, Stefan G Kertesz, Sally Satel, Richard A Lawhern, Kate M Nicholson, Rosemary C Polomano, Owen D Williamson, Heath McAnally, Ming-Chih Kao, Stephan Schug, Robert Twillman, Terri A Lewis, Richard L Stieg, Kate Lorig, Theresa Mallick-Searle, Robert W West, Sarah Gray, Steven R Ariens, , Jennifer Sharpe Potter, Penney Cowan, Chad D Kollas, Danial Laird, Barby Ingle, J Julian Grove, Marian Wilson, Kashelle Lockman, Fiona Hodson, Carol S Palackdharry, Roger B Fillingim, PhD, Jeffrey Fudin, Jennifer Barnhouse, Ajay Manhapra, Steven R Henson, Bruce Singer, PhD, Marie Ljosenvoor, Marlisa Griffith, Jason N Doctor, PhD, Kimeron Hardin, Cathleen London, Jon Mankowski, Andrea Anderson, Linda Ellsworth, Lisa Davis Budzinski, Becky Brandt, Greg Hartley, Debbie Nickels Heck, Mark J Zobrosky, Celeste Cheek, Megan Wilson, Cynthia E Laux, Geralyn Datz, Justin Dunaway, Eileen Schonfeld, RN, Chaplain, Melissa Cady, Thérèse LeDantec-Boswell, Meredith Craigie, John Sturgeon, Pamela Flood, Melita Giummarra, Jessica Whelan, Beverly E Thorn, Richard L Martin, Michael E Schatman, Maurice D Gregory, Joshua Kirz, Patti Robinson, James G Marx, Jessica R Stewart, Phillip S Keck, Scott E Hadland, Jennifer L Murphy, Mark A Lumley, Kathleen S Brown, Michael S Leong, Mechele Fillman, James W Broatch, Aaron Perez, Kristine Watford, Kari Kruska,, Dokyoung Sophia You, Stacy Ogbeide, Amy Kukucka, Susan Lawson, James B Ray, PharmD, T Wade Martin, James B Lakehomer, Anne Burke, Robert I Cohen, Peter Grinspoon, Marc S Rubenstein, Stephani Sutherland, Kristie Walters, Travis Lovejoy, International Stakeholder Community of Pain Experts and Leaders Call for an Urgent Action on Forced Opioid Tapering, *Pain Medicine*, 20, Issue 3, March 2019, Pages 429–433, https://doi.org/10.1093/pm/pny228

Delcher C, Wang Y, Wagenaar AC, Goldberger BA, Cook RL, Maldonado-Molina MM. Prescription and illicit opioid deaths and the prescription drug monitoring program in Florida [Letter] Am J Public Health. 2016;106:e10–1. doi: 10.2105/AJPH.2016.303104.

Demidenko MI, Dobscha SK, Morasco BJ, Meath THA, Ilgen MA, Lovejoy TI. Suicidal ideation and suicidal self-directed violence following clinician-initiated prescription opioid discontinuation among long-term opioid users. *Gen Hosp Psychiatry*. 2017;47:29-35. doi:10.1016/j.genhosppsych.2017.04.011

Denney, J.T., Rogers, R.G., Krueger, P.M. and Wadsworth, T. (2009), Adult Suicide Mortality in the United States: Marital Status, Family Size, Socioeconomic Status, and Differences by Sex*. Social Science Quarterly, 90: 1167-1185. doi:10.1111/j.1540-6237.2009.00652.x

Department of Veteran Affairs. (2015) VA Opioid Prescription Policy, Practice, and Procedures. 114th Congress, 4th Hearing. https://www.govinfo.gov/content/pkg/CHRG-114shrg94295/html/CHRG-114shrg94295.htm

Deyo RA, Hallvik SE, Hildebran C, Marino M, Springer R, Irvine JM, O'Kane N, Van Otterloo J, Wright DA, Leichtling G, Millet LM, Carson J, Wakeland W, McCarty D. Association of Prescription Drug Monitoring Program Use With Opioid Prescribing and Health Outcomes: A Comparison of Program Users and Nonusers. J Pain. 2018 Feb;19(2):166-177. doi: 10.1016/j.jpain.2017.10.001. Epub 2017 Oct 18. PMID: 29054493; PMCID: PMC5803414.

Dubin Ruth, Clarke Hance, and Kahan, Mel. The Risks of Opioid Tapering or Rapid Discontinuation. Canadian Family Physician. 2017. Available at: https://www.cfp.ca/news/2017/12/05/12-05

Edlund, M. J., Martin, B. C., Fan, M. Y., Devries, A., Braden, J. B., & Sullivan, M. D. (2010). Risks for opioid abuse and dependence among recipients of chronic opioid therapy: results from the TROUP study. *Drug and alcohol dependence*, *112*(1-2), 90–98. https://doi.org/10.1016/j.drugalcdep.2010.05.017

Edlund, Mark J., Bradley C. Martin, Joan E. Russo, Andrea DeVries, Jennifer Brennan Braden, and Mark D. Sullivan. 2014. "The Role of Opioid Prescription in Incident Opioid Abuse and Dependence Among Individuals With Chronic Noncancer Pain: the Role of Opioid Prescription," The Clinical Journal of Pain 30(7): 557–564.

Edmonds, A. T., Bensley, K. M., Hawkins, E. J., & Williams, E. C. (2020). Geographic differences in receipt of addictions treatment in a national sample of patients with alcohol use disorders from the U.S. Veterans Health Administration. *Substance abuse*, 1–10.

Finley, E. P., Bollinger, M., Noël, P. H., Amuan, M. E., Copeland, L. A., Pugh, J. A., Dassori, A., Palmer, R., Bryan, C., & Pugh, M. J. (2015). A national cohort study of the association between the polytrauma clinical triad and suicide-related behavior among US Veterans who served in Iraq and Afghanistan. *American journal of public health*, *105*(2), 380–387. https://doi.org/10.2105/AJPH.2014.301957 Fishbain DA. The association of chronic pain and suicide. Seminars in Clinical Neuropsychiatry. 1999 Jul;4(3):221-227. DOI: 10.153/scnp00400221.

Fishbain D. A. (1999). The association of chronic pain and suicide. *Seminars in clinical neuropsychiatry*, *4*(3), 221–227. https://doi.org/10.153/SCNP00400221

Fishbain, D. A., Goldberg, M., Rosomoff, R. S., & Rosomoff, H. (1991). Completed suicide in chronic pain. *The Clinical journal of pain*, *7*(1), 29–36. https://doi.org/10.1097/00002508-199103000-00006

Florence, C. S., Zhou, C., Luo, F., & Xu, L. (2016). The Economic Burden of Prescription Opioid Overdose, Abuse, and Dependence in the United States, 2013. *Medical care*, *54*(10), 901–906. https://doi.org/10.1097/MLR.0000000000000625

Ford JA, Perna D. Prescription drug misuse and suicidal ideation: Findings from the National Survey on Drug Use and Health. *Drug Alcohol Depend.* 2015;157:192-196. doi:10.1016/j.drugalcdep.2015.10.010

García MC, Heilig CM, Lee SH, et al. Opioid prescribing rates in nonmetropolitan and metropolitan counties among primary care providers using an electronic health record system — United States, 2014—2017. CDC Morbidity and Mortality Weekly Report.

Gellad WF, Good CB, Shulkin DJ. Addressing the Opioid Epidemic in the United States: Lessons From the Department of Veterans Affairs. *JAMA Intern Med.* 2017;177(5):611–612. doi:10.1001/jamainternmed.2017.0147

Gironda, R. J., Clark, M. E., Massengale, J. P., & Walker, R. L. (2006). Pain among veterans of Operations Enduring Freedom and Iraqi Freedom. *Pain medicine (Malden, Mass.)*, *7*(4), 339–343. https://doi.org/10.1111/j.1526-4637.2006.00146.x

Good, Bernie C. MD, MPH Federal Healthcare System Experience: VA Implementation, Maintenance, and Evaluation. Veterans Affairs Pharmacy Benefits Management May, 2017.

Government Accountability Office. Progress Made Towards Improving Opioid Safety, but Further Efforts to Assess Progress and Reduce Risk are Needed. 2018. Available at: GAO-18-380, VA HEALTH CARE: Progress Made Towards Improving Opioid Safety, but Further Efforts to Assess Progress and Reducce Risk Are Needed

Hadlandsmyth, K., Mosher, H., Vander Weg, M. W., & Lund, B. C. (2018). Decline in Prescription Opioids Attributable to Decreases in Long-Term Use: A Retrospective Study in the Veterans Health Administration 2010-2016. *Journal of general internal medicine*, *33*(6), 818–824. https://doi.org/10.1007/s11606-017-4283-8

Haffajee, R. L., Mello, M. M., Zhang, F., Zaslavsky, A. M., Larochelle, M. R., & Wharam, J. F. (2018). Four States With Robust Prescription Drug Monitoring Programs Reduced Opioid Dosages. *Health affairs (Project Hope)*, *37*(6), 964–974. https://doi.org/10.1377/hlthaff.2017.1321

Hedegaard H, Curtin SC, Warner M. Suicide mortality in the United States, 1999–2017. NCHS Data Brief, no 330. Hyattsville, MD: National Center for Health Statistics. 2018.

Hooley, J.M., Franklin, J.C. & Nock, M.K. Chronic Pain and Suicide: Understanding the Association. *Curr Pain Headache Rep* 18, 435 (2014). https://doi.org/10.1007/s11916-014-0435-2

Hudson, T. J., Painter, J. T., Martin, B. C., Austen, M. A., Williams, J. S., Fortney, J. C., Sullivan, M. D., & Edlund, M. J. (2017). Pharmacoepidemiologic analyses of opioid use among OEF/OIF/OND veterans. *Pain*, *158*(6), 1039–1045. https://doi.org/10.1097/j.pain.0000000000000874

Ilgen MA, Kleinberg F, Ignacio RV, et al. Noncancer Pain Conditions and Risk of Suicide. *JAMA Psychiatry*. 2013;70(7):692–697. doi:10.1001/jamapsychiatry.2013.908

Johannes, Catherine B., T. Kim Le, Xiaolei Zhou, Joseph A. Johnston, and Robert H. Dworkin. 2010. "The Prevalence of Chronic Pain in United States Adults: Results of an Internet-Based Survey," Journal of Pain 11(11): 230-1239.

Johnson, C.E., Bush, R.L., Harman, J., Bolin, J., Evans Hudnall, G. and Nguyen, A.M. (2015), Variation in Utilization of Health Care Services for Rural VA Enrollees With Mental Health-Related Diagnoses. The Journal of Rural Health, 31: 244-253. doi:10.1111/jrh.12105

Kang, H. K., Bullman, T. A., Smolenski, D. J., Skopp, N. A., Gahm, G. A., & Reger, M. A. (2015). Suicide risk among 1.3 million veterans who were on active duty during the Iraq and Afghanistan wars. *Annals of epidemiology*, *25*(2), 96–100.

Kaplan, Mark S., McFarland, Bentson H., Huguet, Nathalie, and Valenstein, Marcia. 2012: Suicide Risk and Precipitating Circumstances Among Young, Middle-Aged, and Older Male Veterans American Journal of Public Health 102, S131_S137, https://doi.org/10.2105/AJPH.2011.300445

Keyes, K. M., Cerdá, M., Brady, J. E., Havens, J. R., & Galea, S. (2014). Understanding the rural-urban differences in nonmedical prescription opioid use and abuse in the United States. *American journal of public health*, *104*(2), e52–e59. https://doi.org/10.2105/AJPH.2013.301709

Kilby AA. Opioids for the Masses: Welfare Tradeoffs in the Regulation of Narcotic Pain Medications. Cambridge, MA: Massachusetts Institute of Technology; 2015.

Kim HM, Smith EG, Ganoczy D, et al. Predictors of suicide in patient charts among patients with depression in the Veterans Health Administration health system: importance of prescription drug and alcohol abuse. *J Clin Psychiatry*. 2012;73(10):e1269-e1275. doi:10.4088/JCP.12m07658

Kuramoto, S. J., Chilcoat, H. D., Ko, J., & Martins, S. S. (2012). Suicidal ideation and suicide attempt across stages of nonmedical prescription opioid use and presence of prescription opioid disorders among U.S. adults. *Journal of studies on alcohol and drugs*, *73*(2), 178–184. https://doi.org/10.15288/jsad.2012.73.178

Lewei, A.L., Peltzman, T., McCarthy, J.F., et al. (2019). Changing trends in opioid overdose deaths and prescription opioid receipt among veterans. *American Journal of Preventive Medicine,* 57(1), 106-110. https://doi.org/10.1016/j.amepre.2019.01.016

Lin LA, Bohnert ASB, Kerns RD, Clay MA, Ganoczy D, Ilgen MA. Impact of the Opioid Safety Initiative on opioid-related prescribing in veterans. Pain. 2017 May;158(5):833-839. doi: 10.1097/j.pain.0000000000000837. PMID: 28240996.

Lund BC, Ohl ME, Hadlandsmyth K, Mosher HJ. Regional and Rural-Urban Variation in Opioid Prescribing in the Veterans Health Administration. *Mil Med*. 2019;184(11-12):894-900. doi:10.1093/milmed/usz104

Maguen S, Luxton DD, Skopp NA, et al. Killing in combat, mental health symptoms, and suicidal ideation in Iraq war veterans. *J Anxiety Disord*. 2011;25(4):563-567. doi:10.1016/j.janxdis.2011.01.003

McCarthy, John F., Blow, Frederic C., Ignacio, Rosalinda C., Ilgen, Mark A., Austin, Karen L. and Valenstein, Marcia, 2012:Suicide Among Patients in the Veterans Affairs Health System: Rural-Urban Differences in Rates, Risks, and Methods American Journal of Public Health **102**, S111_S117, **https://doi.org/10.2105/AJPH.2011.300463**

Meinhofer A. Prescription drug monitoring programs: the role of asymmetric information on drug availability and abuse. Am J Health Econ. 2017

Mohamed S, Neale M, Rosenheck RA. VA intensive mental health case management in urban and rural areas: veteran characteristics and service delivery. *Psychiatr Serv*. 2009;60(7):914-921. doi:10.1176/ps.2009.60.7.914

Moyo, P., Simoni-Wastila, L., Griffin, B. A., Onukwugha, E., Harrington, D., Alexander, G. C., and Palumbo, F. (2017) Impact of prescription drug monitoring programs (PDMPs) on opioid utilization among Medicare beneficiaries in 10 US States. *Addiction*, 112: 1784– 1796. doi: 10.1111/add.13860.

National Center for Veterans Analysis and Statistics (NCVAS). VA Utilization Profile FY 2017. 2017. https://www.va.gov/vetdata/docs/Quickfacts/VA_Utilization_Profile_2017.pdf

Oliva Elizabeth M, Bowe Thomas, Manhapra Ajay, Kertesz Stefan, Hah Jennifer M, Henderson Patricia et al. Associations between stopping prescriptions for opioids, length of opioid treatment, and overdose or suicide deaths in US veterans: observational evaluation BMJ 2020; 368 :m283

Oquendo, M. A., & Volkow, N. D. (2018). Suicide: A Silent Contributor to Opioid-Overdose Deaths. *New England Journal of Medicine*, *378*(17), 1567–1569. https://doi.org/10.1056/NEJMp1801417

Park, Tae Woo, Richard Saitz, Dara Ganoczy, Mark A. Ilgen, and Amy S.B. Bohnert. 2015. "Benzodiazepine Prescribing Patterns and Deaths from Drug Overdose Among U.S. Veterans Receiving Opioid Analgesics: Case-Cohort Study," bmj 350: h2698.

Petzel, Robert., Written statement of Dr. Robert Petzel Under Secretary for Health Veterans Health Administration Department Of Veterans Affairs Before The Senate Veterans' Affairs Committee U.S. Senate April 30, 2014

Possemato K, Wray LO, Johnson E, Webster B, Beehler GP. Facilitators and Barriers to Seeking Mental Health Care Among Primary Care Veterans With Posttraumatic Stress Disorder. *J Trauma Stress*. 2018;31(5):742-752. doi:10.1002/jts.22327

Sher, L., & Yehuda, R. (2011). Preventing suicide among returning combat veterans: A moral imperative. *Military Medicine, 176*(6), 601–602. https://doi.org/10.7205/MILMED-D-11-00026

J. M. C. Santos Silva & Silvana Tenreyro, 2006. "The Log of Gravity," The Review of Economics and Statistics, MIT Press, vol. 88(4), pages 641-658, November.

Sandbrink, F, 2019. "Session II Efforts to reduce high-dose opioid analgesic prescribing: What have we learned?", Pages 1-33.

Smalley KB, Yancey CT, Warren JC, Naufel K, Ryan R, Pugh JL. Rural mental health and psychological treatment: a review for practitioners. J Clin Psychol. 2010;66(5):479-489.

Suffoletto B, Lynch M, Pacella CB, Yealy DM, Callaway CW. The Effect of a Statewide Mandatory Prescription Drug Monitoring Program on Opioid Prescribing by Emergency Medicine Providers Across 15 Hospitals in a Single Health System. J Pain. 2018 Apr;19(4):430-438. doi: 10.1016/j.jpain.2017.11.010. Epub 2017 Dec 11. PMID: 29241835; PMCID: PMC5869080.

Sun Eric C, Dixit Anjali, Humphreys Keith, Darnall Beth D, Baker Laurence C, Mackey Sean et al. Association between concurrent use of prescription opioids and benzodiazepines and overdose: retrospective analysis BMJ 2017; 356 :j760

Swensen, Isaac D, Substance-abuse treatment and mortality, Journal of Public Economics, Volume 122, 2015, Pages 13-30, ISSN 0047-2727, https://doi.org/10.1016/j.jpubeco.2014.12.008.

Talari MP, Yarra P Increased risk of death by suicide or overdose in patients stopping opioid medications *Evidence-Based Nursing* Published Online First: 31 August 2020. doi: 10.1136/ebnurs-2020-103293

Tang, Nicole K. Y., and Catherine Crane. "Suicidality in Chronic Pain: a Review of the Prevalence, Risk Factors and Psychological Links." Psychological Medicine, vol. 36, no. 5, 2006, pp. 575–586., doi:10.1017/S0033291705006859.

Teich, J., Ali, M.M., Lynch, S. and Mutter, R. (2017), Utilization of Mental Health Services by Veterans Living in Rural Areas. The Journal of Rural Health, 33: 297-304. doi:10.1111/jrh.12221

Toblin, Robin L., Phillip J. Quartana, Lyndon A. Riviere, Kristina Clarke Walper, and Charles W. Hoge. 2014. "Chronic Pain and Opioid Use in U.S. Soldiers After Combat Deployment," Journal of the American Medical Association: Internal Medicine 174(8): 1400- 1401.

Tondo, L., Albert, M. J., & Baldessarini, R. J. (2006). Suicide Rates in Relation to Health Care Access in the United States: An Ecological Study. *The Journal of Clinical Psychiatry, 67*(4), 517–523. https://doi.org/10.4088/JCP.v67n0402

United States Department of Agriculture Economic Research Service. Measuring rurality: Rural-urban continuum codes, 2013. Available at*:* https://www.ers.usda.gov/data-products/rural-urban-continuum-codes.aspx

Van Zee A. The Promotion and Marketing of OxyContin: Commercial Triumph, Public Health Tragedy. *Am J Public Health*. 2009;99(2):221-227. doi:10.2105/AJPH.2007.131714.

Vanderlip, E. R., Sullivan, M. D., Edlund, M. J., Martin, B. C., Fortney, J., Austen, M., Williams, J. S., & Hudson, T. (2014). National study of discontinuation of long-term opioid therapy among veterans. *Pain*, *155*(12), 2673–2679. https://doi.org/10.1016/j.pain.2014.09.034

Wallace AE, Weeks WB, Wang S, Lee AF, Kazis LE. Rural and urban disparities in health-related quality of life among veterans with psychiatric disorders. Psychiatr Serv. 2006;57(6):851-856.

Wenger, Jennie W., Caolionn O'Connell, and Linda Cottrell, Examination of Recent Deployment Experience Across the Services and Components. Santa Monica, CA: RAND Corporation, 2018. https://www.rand.org/pubs/research_reports/RR1928.html.

Wilcox, H. C., K. R. Conner, and E. D. Caine. 2004. Association of alcohol and drug use disorders and completed suicide: an empirical review of cohort studies. Drug and Alcohol Dependence 76:S11–19.

Wide-ranging online data for epidemiologic research (WONDER), Atlanta, GA: CDC, National Center for Health Statistics; 2020. Available at http://wonder.cdc.gov




Figure 1: Quarterly evolution of unique VA patient opioid prescriptions (source: Good, 2017)




Figure 2: Timeline of VA opioid policies and OSI implementation (source: Sandbrink, 2019)



Figure 3 Average county-year suicide trend for veteran/military personnel and civilians




Figure 4: Coefficients from a linear regression of suicides per 100,000 among veterans/military personnel and civilians by urban-rural status. The specification includes right-hand side controls discussed in the text.

Table 1: Descriptive Statistics – Pre- OSI Suicides, Economic Conditions & Addiction Treatment

Variable Description	Veteran		Civilian	
(county-year)	**Rural**	**Urban**	**Rural**	**Urban**
Suicide count	0.71	3.73	2.94	14.69
	(1.13)	(5.20)	(3.26)	(20.79)
Population estimates	2,127	12,761	17,291	115,051
	(1,919)	(15,138)	(14,488)	(156,550)
Suicide rate per 100,000	34.58	30.20	18.34	14.39
	(66.26)	(32.43)	(21.57)	(10.21)
Median income	$30,437	$39,095	$20,541	$25,995
	($7,783)	($10,411)	($4,741)	($6,562)
Unemployment rate	7.94%	7.04%	9.07%	8.35%
	(6.74%)	(4.02%)	(3.94%)	(2.70%)
Percent of population	5.66%	7.50%	55.68%	57.39%
that is female	(3.48)	(3.55%)	(4.31%)	(2.80%)
Percent of population	6.16%	7.86%	28.49%	31.25%
that is 18-34 years old	(4.16%)	(4.31%)	(5.91%)	(6.85%)
Percent of population	42.86%	38.03%	17.91%	14.74%
that is 65 year and older	(9.57%)	(8.68%)	(4.18%)	(3.33%)
Percent of population	17.60%	24.41%	16.52%	24.86%
that has a bachelors	(9.00%)	(11.00%)	(7.61%)	(11.46%)

	Rural	**Urban**
Addiction treatment	1.44	8.76
centers	(1.98)	(15.64)
Addiction treatment	7.48	5.64
centers per 100,000	(10.85)	(6.21)

Note: Standard deviations for each variable are in parentheses below the mean.

Table 2: Descriptive Statistics – Pre & Post OSI Implementation

Variable Description	**Veteran**		**Civilian**	
(county-year)	**Pre-OSI**	**Post-OSI**	**Pre-OSI**	**Post-OSI**
Suicide count	2.00	2.19	7.97	9.48
	(3.81)	(3.95)	(14.99)	(17.05)
Population estimates	6,679	6,288	59,138	63,064
	(11,308)	(10,610	(113792)	(121604)
Suicide rate per 100,000	32.70	43.44	16.65	18.52
	(54.46)	(76.81)	(17.73)	(16.26)
Median income	$34,143	$34,724	$22,876	$23,593
	($9,969)	($10,344)	($6,210)	($6,161)
Unemployment rate	7.55%	7.86%	8.76%	8.93%
	(5.75)	(6.15)	(3.48)	(3.53)
Percent of population that is female	6.5%	6.8%	56.4%	55.6%
	(3.63)	(3.75)	(3.83)	(3.67)
Percent of population that is 18-34 years old	6.9%	7.1%	29.7%	29.3%
	(4.30)	(4.51)	(6.48)	(6.27)
Percent of population that is 65 year and older	40.8%	46.4%	16.6%	18%
	(9.51)	(10.26)	(4.14)	(4.42)
Percent of population that has a bachelors	20.5%	21.2%	20.1%	21.1%
	(10.46)	(10.75)	(10.31)	(10.57)
	Pre-OSI		**Post-OSI**	
Addiction treatment centers	4.57		5.19	
	(10.96)		(12.04)	
Addiction treatment centers per 100,000	6.7		7.27	
	(9.20)		(9.56)	

Note: Standard deviations for each variable are in parentheses below the mean.

Table 3: OSI Regression Results (OLS)

	OLS (without controls)	OLS (with controls)
Dependent Variable	**Suicides/100k**	**Suicides/100k**
Variables		
Civilian Suicide Rate by Year (base: 2010)		
2011	0.601**	0.552
	(0.270)	(0.519)
2012	-0.715*	-1.072
	(0.383)	(0.876)
2013	-0.081	-0.615
	(0.350)	(1.049)
2014	0.686*	-0.401
	(0.351)	(1.188)
2015	2.294***	0.729
	(0.375)	(1.284)
2016	2.588***	0.467
	(0.393)	(1.523)
Veteran*Year (base: 2010)		
2011	0.762	1.549
	(1.052)	(1.143)
2012	1.515	2.047
	(1.074)	(1.329)
2013	1.091	1.591
	(1.076)	(1.371)
2014	1.853	2.007
	(1.293)	(1.543)
2015	3.797***	3.390**
	(1.264)	(1.541)
2016	4.752***	3.932***
	(1.238)	(1.434)
Year*Rural (base: 2010)		
2011	0.978	-0.130
	(0.718)	(1.056)
2012	0.878	-1.551
	(0.805)	(1.522)
2013	1.358*	-2.018
	(0.814)	(1.890)
2014	0.764	-2.774
	(0.805)	(2.110)
2015	2.000**	-0.495
	(0.792)	(2.209)
2016	1.282	-0.131
	(0.823)	(2.542)

DDD: Veteran*Year*Rural (base: 2010)		
2011	**2.472**	**2.029**
	(2.698)	**(2.834)**
2012	**2.841**	**3.290**
	(2.853)	**(3.194)**
2013	**10.122*** **	**11.099*** **
	(2.955)	**(3.381)**
2014	**8.941*** **	**9.598*** **
	(2.991)	**(3.508)**
2015	**8.341*** **	**9.723*** **
	(3.046)	**(3.516)**
2016	**8.191*** **	**10.279*** **
	(3.075)	**(3.556)**

Note: *p<0.1, **p<0.05, ***p<0.01. The dependent variable in the linear model is the suicide rate per 100,000 individuals at the county-year level. Both regressions are weighted by the total population in each county and include county and year fixed effects with standard errors clustered by county. Control variables in column 2 include: PDMP indicators, number of treatment centers, interactions between PDMP policies and treatment centers, and estimates for the percentage of the population aged 18-34, female, and black, the unemployment rate, median income, and percentage that have some college experience, all for both veteran and non-veteran populations separately. Finally, all control variables are interacted with veteran and rural county status.

Table 4: OSI Regression Results (Poisson)

	Poisson QMLE (without controls)	Poisson QMLE (with controls)
Dependent Variable	**Suicide Count**	**Suicide Count**
Variables		
Civilian Suicide Rate by Year (base: 2010)		
2011	0.015	0.005
	(0.024)	(0.064)
2012	-0.070	-0.055
	(0.048)	(0.093)
2013	-0.014	0.051
	(0.030)	(0.090)
2014	0.057*	0.141
	(0.032)	(0.111)
2015	0.171***	0.278*
	(0.063)	(0.148)
2016	0.188***	0.297*
	(0.068)	(0.171)
Veteran*Year (base: 2010)		
2011	0.007	0.044
	(0.049)	(0.054)
2012	0.089*	0.152***
	(0.050)	(0.056)
2013	0.039	0.105*
	(0.055)	(0.061)
2014	0.004	0.086
	(0.076)	(0.081)
2015	-0.006	0.071
	(0.085)	(0.092)
2016	0.010	0.083
	(0.090)	(0.099)
Year*Rural (base: 2010)		
2011	-0.007	-0.036
	(0.048)	(0.078)
2012	0.046	-0.066
	(0.069)	(0.110)
2013	0.016	-0.167
	(0.060)	(0.115)
2014	-0.048	-0.283**
	(0.057)	(0.133)
2015	0.005	-0.253
	(0.077)	(0.164)
2016	-0.057	-0.315*
	(0.084)	(0.185)

DDD: Veteran*Year*Rural (base: 2010)		
2011	**0.191***	**0.185**
	(0.112)	**(0.116)**
2012	**0.148**	**0.186**
	(0.114)	**(0.126)**
2013	**0.322****	**0.388*****
	(0.126)	**(0.139)**
2014	**0.369*****	**0.438*****
	(0.133)	**(0.145)**
2015	**0.324****	**0.411*****
	(0.128)	**(0.139)**
2016	**0.306****	**0.408*****
	(0.132)	**(0.143)**

Note: *p<0.1, **p<0.05, ***p<0.01. The dependent variable in the Poisson QMLE specification is the count of suicides at the county-year level. Both regressions are weighted by the total population in each county and include county and year fixed effects with standard errors clustered by county. Control variables in column 2 include: PDMP indicators, number of treatment centers, interactions between PDMP policies and treatment centers, and estimates for the percentage of the population aged 18-34, female, and black, the unemployment rate, median income, and percentage that have some college experience, all for both veteran and non-veteran populations separately. Finally, all control variables are interacted with veteran and rural county status.

Table 5: Veteran Migration Regression Results

	OLS
Dependent Variable	**Percent Veteran**
Variables	
Year (base: 2010)	
2011	-0.003***
	(0.000)
2012	-0.005***
	(0.000)
2013	-0.009***
	(0.000)
2014	-0.012***
	(0.000)
2015	-0.015***
	(0.000)
2016	-0.018***
	(0.000)
Rural*Year (base: 2010)	
2011	0.000
	(0.000)
2012	0.000
	(0.000)
2013	0.000
	(0.000)
2014	0.000
	(0.000)
2015	0.000
	(0.000)
2016	0.001
	(0.000)

Note: The dependent variable is the percent of a county's population that is made up of veterans and military personnel; this regression is weighted by total county population and includes year and county fixed effects. *p<0.1, **p<0.05, ***p<0.01