COVER CROPS, DROUGHT, YIELD AND RISK: AN ANALYSIS OF US. SOYBEAN PRODUCTION

Fengxia Dong
USDA-ERS
Fengxia.dong@usda.gov

The findings and conclusions in this paper are those of the author(s) and should not be construed to represent any official USDA or U.S. Government determination or policy.

Abstract

Besides a variety of production and environmental benefits, cover cropping has been advocated as a means to increase resilience to drought. We explored factors that affected farmer's adoption of cover crops and examined the effects of cover crops on soybean yield and its risk using USDA's 2018 ARMS Phase II Soybean Production Practices and Costs Report and Phase III Soybean Costs and Returns Report. Incorporating drought occurrence in current year and previous 5 years into our analysis, we find that previous occurrence of drought did not affect farmers' adoption of cover crops and the effects of cover crops on yield and its risk are mixed. Under a drought condition, cover crops reduced soybean yield and increased yield variation; but in the meantime, they also reduced the risk of crop failure, or made yield less negatively skewed. The insignificant effect of previous drought on cover crop adoption and the mixture of positive and negative effects of cover crops on yield and its risk imply that farmers are divided in their acceptance of cover crops as a means to build resilience to drought.

Introduction

Climate change has caused increasing frequency and severity of drought stress in the U.S. Water scarcity has become one of the most severe constraints to agricultural production, adversely affecting crop yields and presenting a major challenge to sustainable food production. Along with prescribed grazing, mulching, micro-irrigation, and conservation tillage, cover crop is among the five short- and long-term strategies for dealing with drought conditions that the Natural Resources Conservation Service (NRCS) financially assists farmers with (USDA Climate Hubs 2021). Cover crops can be broadly defined as any non-cash crop grown in addition to the primary cash crop (UC Sustainable Agriculture Research and Education (SARE) 2017). The use of cover crops as a cropping strategy is not new. It had been practiced by people in ancient Greece, Rome, and China as early as 3,000 years ago (Langdale et al. 1991). Cover crops were first used in the 18th century in the U.S. and extensively expanded in the 19th century (Groff 2015), although their role by the time mainly as green manures. The affordability and ease of use of synthetic fertilizer at the end of World War II, however, had attracted farmers to utilize more synthetic fertilizer instead of cover crops to further improve crop yields. The use of cover crops in conventional agriculture has gradually become less common since then (Groff 2015).

Cover crops can protect and improve soil between periods of regular crop production (Schnepf and Cox 2006). Besides a variety of production, soil health, and environmental benefits such as increasing weed and pest suppression, reducing runoff of sediments and nutrients into waterways, and reducing soil erosion and compaction, cover crops can improve water infiltration, reduce water evaporation, and increase soil's water holding capacity (e.g., USDA Natural Resources Conservation Services (NRCS) 2018; Mitchell et al. 2015; Blanco-Canqui et al. 2015; McDaniel, Tiemann, and Grandy 2014; Laloy and Bielders 2010; Dean and Weil 2009;

Sainju et al. 2002, 2006). Given financial support, cover crop acres enrolled in the Environmental Quality Incentives Program (EQIP) increased from 312.6 thousand acres in 2009 to 2,443.1 thousand acres in 2020 (Climate Hubs, USDA, 2021). In addition, recently, farmers have been paid to plant cover crops by large seeds and chemical companies to generate carbon offset credits for the seeds and chemicals makers (Reuters 2022).

However, the adoption cover crop has been slow. There are well recognized tradeoffs and limitations in adopting cover crop as a conservation strategy (SARE 2017). Besides costs of soil preparation, seeds, and labor, there are also challenges in implementation and management, such as cover crop species, planting and termination time which may interfere with fall harvest or spring planting, and producing too much surface residue (CTIC 2015; Sackett 2013; Miller, Chin, and Zook 2012; Snapp et al. 2005). Moreover, there are concerns that water need by cover crop may reduce the amount of water available to the following main crop (SARE 2017; Clark et al. 1997; Corak et al., 1991; Ebelhar et al., 1984; Munawar et al., 1990). Unlike conservation tillage having become well adopted in 67% of soybean fields, only about 8.4 percent of soybean fields were planted cover crops in 2018 (NASS 2019).

Along with above concerns, effect on crop yield and risk is another important factor in farmer's adoption decision of cover crops as yield and risk directly affect farmer's economic returns. The Iowa Farm and Rural Life Poll in 2015 (Arbuckle 2016) reveals that 74% of the farmers consider that economic factors have a moderate to very strong influence on their changes in management practices. The 2017 Cover Crop Survey conducted by SARE (2017) also shows that fear of a lack of economic returns (54% of respondents), increasing production risk (48% of respondents), and potential yield reduction (44%) are among the major concerns for non-users. Another cover crop survey conducted in 2015 shows that potential yield benefits to cash crops

was an important factor in decision making, especially for non-adopters (CTIC 2015). Therefore, analysis of yield and its risk of management practices is essential to overcome the barrier by setting up effective supporting programs to promote good management practice adoption.

The results of existing studies of yield effects of cover crops are mixed. A meta-analysis of the response of corn yield to cover crops by Miguez and Bollero (2005) concludes that legume cover crops increase corn yield by 37%. Similarly, Andraski and Bundy (2005) and Munoz et al. (2014) find a positive effect of cover crop biomass on corn yields. Contrastingly, Reddy (2017) finds lower soybean yield with cover crops compared with no cover crops. Nielsen et al. (2016) find that there was an average 10% reduction in wheat yield following a cover crop compared with following fallow, regardless of whether the cover crop was grown in a mixture or in a single-species planting. In addition, yield reductions were greater under drier conditions. And Smith et al. (2014), Hunter et. al (2019), and Acuna and Villamil (2014) find no benefits of growing cover crop on subsequent soybean or other crop yield.

Previous studies of yield effects of cover crops are mainly conducted in field experimental plots using agronomic models. A study based on a large number of fields with different agroecological characteristics and under varied weather is in need. Increasing frequency and severity of adverse events can expose farms to significant production uncertainty. Therefore, special attention is paid to downside risk exposure. In general, downside risk is the risk that is associated with unfavorable events and located in the lower tail of the yield or return distribution (Kim et al. 2014). As pointed out by Hardaker et al. (2004), Kim et al. (2014), and OECD (2011), analyzing both the exposure to risk and levels of downside risk in agriculture is a key component in assessing welfare impacts. While cover cropping is recommended to farmers to

deal with drought, its effects on farm yield risk and especially downside risk are not well documented. This study aims to fill the literature gap.

The paper makes three contributions. First, it explores the factors that affect the adoption of cover cropping. We consider not only land characteristics and farmer's concerns and demographics, but also droughts in previous years. This enables us to reveal whether farmers view cover cropping as an effective means to increase resilience to drought. Second, the paper examines the effects of cover crops on yield variation and downside risk. We employ moments of yield distribution to evaluate the exposure to yield variation and downside risk. Disentangling the productivity effects of adaptation is of paramount importance. It will reveal whether farmers who adopt cover crop are indeed getting benefits in terms of an increase in crop productivity, a benefit crucial to a broader adoption. Third, the paper utilizes a data set covering the majority of soybean fields in the U.S. with significantly different soil types and weather conditions. Two interplays, one between soil types and the adaptation strategy (namely, cover cropping), the other between weather and the cover cropping practice, are included in yield and its risk analysis. The interplays along with farmer demographics, farm characteristics, and input use allow us to examine effectiveness of the managerial options for risk mitigation, especially in the threat of drought.

The rest of the paper is organized as follows. Next section discusses the empirical framework, followed by a description of data and variables. Then estimation results are discussed, followed by conclusions at the end.

Empirical Framework

To investigate the impacts of cover crops on crop yield and yield risks, we start with the moment functions of crop yield.

Moment Representation of Production Function

Our empirical model relies on central-moment measurements. It is based on Antle's (1983) moment-based approach, which provides a flexible and convenient basis for evaluating exposure to production risk. As discussed in Antle (1983), a stochastic production function can be represented by a general parameterization of the moment functions. Consider a farmer who uses a vector of inputs $\boldsymbol{x}$ and drought adaptation strategies (such as cover cropping) C to produce a scalar output Q. Using $\boldsymbol{\beta_1}$ to indicate a vector of technology parameters and $\mathbf{e}$ uncontrollable random factors such as weather, droughts, pests, and diseases, the production technology can be represented by $Q = Q(\boldsymbol{x}, C, \boldsymbol{e}, \boldsymbol{\beta_1})$. The production function is stochastic given the random error term $\mathbf{e}$. Let the stochastic output Q have a cumulative distribution $F(\boldsymbol{e})$, then the first and the ith central moments of output Q can be represented, respectively, as

$$m_1(\boldsymbol{x}, C, \boldsymbol{\beta_1}) = E[Q(\boldsymbol{x}, C, \boldsymbol{e}, \boldsymbol{\beta_1})] = \int Q(\boldsymbol{x}, C, \boldsymbol{e}, \boldsymbol{\beta_1}) dF(\boldsymbol{e}) \quad (1)$$

$$m_i(\boldsymbol{x}, C, \boldsymbol{\beta_i}) = E[\{Q(\boldsymbol{x}, C, \boldsymbol{e}, \boldsymbol{\beta_1}) - m_1(\boldsymbol{x}, C, \boldsymbol{\beta_1})\}^i]$$

$$= \int (Q(\boldsymbol{x}, C, \boldsymbol{e}, \boldsymbol{\beta_1}) - m_1(\boldsymbol{x}, S, \boldsymbol{\beta_1}))^i dF(\boldsymbol{e}) \qquad for\ i \geq 2 \quad (2)$$

Here E is the expectation operator; m_1 is the mean and m_i is the ith moment of output (for $i \geq 2$); and $\boldsymbol{\beta_i}$ is a vector of parameters. The models in (1) and (2) have the advantage of being flexible as there are no restrictions within or cross moments.

By rewriting equations (1) and (2), we get the following equations

$$\varepsilon_1 = Q(\boldsymbol{x}, S, \boldsymbol{\beta_1}) - m_1(\boldsymbol{x}, S, \boldsymbol{\beta}_1) \quad (3)$$

$$(\varepsilon_1)^i = m_i(\boldsymbol{x}, S, \boldsymbol{\beta}_i) + \varepsilon_i, \quad i \geq 2. \tag{4}$$

Here, $E(\varepsilon_i) = 0$ and $E(\varepsilon_j \varepsilon_{j'}) = 0$ (j=1, 2, …, n and $j \neq j'$).

As discussed by Kendall and Stuart (1977) and shown in many empirical analyses (e.g., Day 1965, Di Falco and Chavas 2009, Di Falco and Veronesi 2014, Tack, Harri, and Coble 2012, and Anderson, Dillon, and Hardaker 1980), the first three moments including location (mean), dispersion (variance), and skewness (the third moment) of a distribution can adequately approximate a distribution. We therefore choose the first three moments to represent the distribution of yield in our analysis. While the variance (m_2) is a traditional measure of risk, the skewness of output (m_3) measure captures the tail asymmetry of a yield distribution around its mean. A negative (positive) skewness implies a distribution skewed to the left (right). A lower skewness presents a greater exposure to downside risk of unexpected low yield, i.e., crop failure.

From equations (3) and (4), we have the mean, variance, and skewness of yield as the following,

$$Q = m_1(\boldsymbol{x}, C, \boldsymbol{\beta}_1) + \varepsilon_1 \tag{5}$$

$$(\varepsilon_1)^2 = m_2(\boldsymbol{x}, C, \boldsymbol{\beta}_i) + \varepsilon_2, \tag{6}$$

$$(\varepsilon_1)^3 = m_3(\boldsymbol{x}, C, \boldsymbol{\beta}_i) + \varepsilon_3. \tag{7}$$

Here, $E(\varepsilon_j) = 0$ and $E(\varepsilon_j \varepsilon_{j'}) = 0$ ($j = 1,2,3$ and $j \neq j'$). Empirically, if $\boldsymbol{\beta}_1^*$ is a consistent estimator of $\boldsymbol{\beta}_1$ from a sample of observed outputs, then $\varepsilon_1^* = Q(\boldsymbol{x}, C, \boldsymbol{\beta_1}) - m_1(\boldsymbol{x}, C, \boldsymbol{\beta}_1^*)$ is a consistent estimator of $\varepsilon_1 = Q(\boldsymbol{x}, C, \boldsymbol{\beta_1}) - m_1(\boldsymbol{x}, C, \boldsymbol{\beta}_1)$. It also suggests that $(\varepsilon_1^*)^i = m_i(\boldsymbol{x}, S, \boldsymbol{\beta}_i^*) + \varepsilon_i$, $i \geq 2$ is a consistent estimator of $(\varepsilon_1)^i$. The models in (5), (6), and (7) have no restrictions within or cross moments and thus are also flexible.

The adoption of cover crops C is variance increasing, variance neutral, or variance decreasing if $\frac{\partial m_2}{\partial C} > 0, = 0$, or < 0, respectively. For a risk averse farmer, $\frac{\partial m_2}{\partial C} > 0$ meaning that the adoption of cover crops creates greater risk in output is undesirable. Similarly, he adoption of cover crops C is skewness increasing, skewness neutral, or skewness decreasing if $\frac{\partial m_3}{\partial C} > 0, = 0$, or < 0, respectively. And for a risk averse farmer, $\frac{\partial m_3}{\partial C} < 0$ meaning that the adoption of cover crops increasing exposure to a lower output is undesirable.

Some factors known to farmers but unknown to economists may affect both yield and the cover cropping decision. Consequently, when empirically estimating the yield equation as shown in (5), a concern that the adoption of cover crops may be endogenous arises. The endogeneity may result in inconsistent and biased estimates. To address the potential endogeneity issue, we use a two-stage method, which is one of the most potent and versatile tools available to treat endogeneity (Antonakis et al. 2014). We will first estimate the use of cover crops by a logit model with an instrumental variable approach.

We model the adoption of cover crop in a logit model as follows:

Cover crop: $$y_{cc}^{*} = \alpha' \boldsymbol{x} + \gamma Z + \epsilon$$

$$y_{cc} = \begin{cases} 1 & \text{if } y_{cc}^{*} > 0 \\ 0 & \text{otherwise} \end{cases} \quad (8)$$

Here y_{cc}^{*} is a latent continuous variable associated with the adoption of cover cropping; y_{cc} is the corresponding observed binary outcome; β' is a transpose vector for parameters to be estimated; and Z is a vector of instrumental variables.

Data and Statistics

We apply our analysis to U.S. soybean production. Farm-level survey data provide us a good opportunity to look more closely at farm activities and motives for them (Dong, Hennessy, and Jensen 2010; Dong, Hennessy, Jensen, and Volpe 2016). We construct the data from the USDA's 2018 ARMS Phase II, Soybean Production Practices and Costs Report, and Phase III Soybean Costs and Returns Report. The phase II survey covers a cross-section soybean fields in 19 states and collects information on production and management practices, input uses, and field characteristics. The phase III report provides information on farm operator and financial characteristics. As conventional and organic production are significantly different in production practices, we only use data from conventional soybean growers.

We delete all observations with missing values, leaving a total of 1,177 observations. ARMS has a complex survey design and is a probability-based survey with unequal probability sampling (National Research Council, 2008). To account for the survey design, we use the sampling weights (expansion factors) provided by NASS to expand the sample to generate population estimates in the statistical analysis. Survey population and sample summary statistics for variables used in the models are reported in Table 1. With the survey weights applied to the sample observations, the weighted sample represents approximately 835,530 soybean fields in the United States.

Variables in Moment Equations

The dependent variables in equations (5) to (7) are the first three moments of the distribution of soybean yield per acre, respectively. Variable *phosphorus* measuring the use of phosphorus per acre is included along with its quadratic term. They are expected to increase the yield. Other independent variables include those on production practices, field characteristics, regional location, and weather/climate. To capture regional differences, indicator variables were

constructed based on Farm Resource Regions (USDA-ERS 2000), which are defined based on farm, soil and climate characteristics rather than state boundaries. The regional dummy variables (*Eastern Uplands, Heartland, Mississippi Portal, Northern Crescent, Northern Great Plains, Prairie Gateway, and Southern Seaboard*) are equal to 1 if the field locates in the corresponding region and 0 otherwise. Two variables (*PlantLate* and *ReplantPct*) are used to capture the impacts of adverse factors negatively affecting soybean yields at the start of the planting season. Variable *PlantLate* is equal to 1 if the planting date fell in the last 15% percentile of the state, and 0 otherwise. Variable *ReplantPct* is the proportion of fields that was replanted. Both might be resulted from adverse weather and result in shorter growing seasons for soybeans, and thus are expected to have a negative effect on yield. Dummy variable *manure* has value 1 if manure was applied to the field and 0 otherwise. Several field characteristics which may affect yield such as soil texture and slope of the field (Butcher et al. 2018; Arora et al. 2011; Shane and Barker 1986; Kaspar et al. 2004; Jiang and Thelen 2004; Kravchenko, Bullock, and Boast 2000; Linkemer Board, and Musgrave 1998; Nelson and Meinhardt 2011) are also included in the yield moment functions. Soil texture is categorized into 5 types: loam, clay, sandy, mixed, and silty. Slope is categorized into 2 levels: nearly level and moderate/steep grade (even or variable). While soybean growing season is from May to September, weather in both July and August are important for soybean yields (Westcott and Jewison 2013). The county-level U.S. Drought Monitor (USDM) indicator jointly produced by the National Drought Mitigation Center (NDMC), the National Oceanic and Atmospheric Administration (NOAA), and the U.S. Department of Agriculture is used. The USDM indicator has five categories: D0-D4, of which D0 indicates abnormally dry but not in drought while D1-D4 indicate moderate drought to exceptional drought. The USDM indicator is based on inputs including the Palmer Drought

Severity Index (PDSI), the Standardized Precipitation Index (SPI), satellite-based assessments of vegetation health, and various indicators of soil moisture as well as hydrologic data (NDMC 2021). If on weekly average over 10% of total county area is categorized as D1 or above in either July or August, then an indicator variable *drought18* is equal to 1 otherwise 0. Moreover, modified growing degree days (mGDD) and overheating growing days (ODD) for soybeans during July and August of 2018 are also included to explicitly capture temperature or heat effects on growth and development of soybean plants. GDD is one of the most important factors influencing the rate of development in soybean (Major et al. 1975; Pedersen and Licht 2014; Kessler, Archontoulis and Lich 2020). Daily mGDD is calculated as

$$mGDD = \max\left(\frac{min(daily\ max\ temp, higher\ development\ threshold) + max(daily\ min\ temp, lower\ development\ thresholds)}{2} - 50, 0\right).$$

GDD depends upon the minimum and maximum temperatures which affect the plant's growth. The higher and lower development thresholds are 86 degrees Fahrenheit and 50 degrees Fahrenheit, respectively, for soybeans. Daily mGDD in July and August is accumulated to get mGDD for the two months. ODD is the count of days in July and August with temperature over 89.6 degrees Fahrenheit and measures the heat stress for crops. The drought adaptation strategy, cover cropping is included in the function as a dummy variable. It is set equal to 1 if the cover cropping was adopted and 0 otherwise. In addition, interactions between cover cropping and drought status *drought18* are also included to capture the effects of the adaptation strategy on yield moments conditional on drought events.

Variables in Logit Model

The instrumental variables used in the cover crop equation include farmers' concerns on soil and water related issues. Seven dummy variables are constructed, indicating concerns on water-drive erosion, wind-driven erosion, soil compaction, poor drainage, low organic matter, water quality, and other concerns, respectively and taking value of 1 if farmer had such a concern and 0 otherwise. We believe that these variables affected cover crop adoption decision but not yield moments. Variable of land ownership is included as another instrumental variable for the same reason. The variable takes value of 1 if the operator owned the land and 0 otherwise. We expect that land ownership may increase the likelihood of fields to adopt cover crops as landowners may care more about soil erosion and soil quality in their own land and thus had more motivation to adopt conservation practices.

Farm size and some other field characteristics such as field size and whether any part of the field was classified as "highly erodible", and whether the field contained a wetland are also included in the cover crop equation given their possible effect on farmer's cover crop decision making (Ding, Schoengold, and Tadesse 2009; Vitale et al. 2011; Wandel and Smithers 2000). Highly erodible land is any land that can erode at excessive rate due to its soil properties. Farmers are required to farm such land in accordance with a conservation plan or system approved by NRCS (USDA Risk Management Agency 2015). We expect that field classified as "highly erodible" is more likely to adopt cover crops for its vulnerability of soil erosion. Both farm and field size are measured in acres. We expect that larger farms and larger fields may be less likely to adopt cover crops given the time and labor requirements of cover crop implementation and management.

Moreover, farmer's age, education, years of experience in farming, and off-farm work are also included. We expect that older farmers or farmers with off-farm work may less likely adopt cover crops given the time and labor investment needed by cover crop implementation and management. Several studies have found evidence that extreme weather affects farmer's adoption of practices. We use variable *drought5yr* to indicate the number of years in which a D1 degree or above drought happened in at least 10% of the county areas during July and August in the last five years. To avoid forbidden regression, all exogenous variables in yield moment equations are also included in the estimation of the cover crop equation. However, since many of them do not have realistic meaning (e.g., phosphorus use should not affect cover crop adoption), we only report several of their estimates, including regional dummy variables, soil texture, and slope of the field in the next section.

Estimation Results

Given the potential endogeneity of cover cropping, we utilize the two-stage-least-square (2SLS) method. In the first stage, we estimate the logit model specified in equation (8) and explore the determinants of the probability of adoption of cover crops with a focus on the effects of climate change, farmer's demographic information and field's characteristics. In the second stage, we estimate the moment equations of (5), (6), and (7).

Results of Logit Model for Cover Crop Adoption

The logit model estimation results are reported in Table 3. The results show that there existed regional difference in the adoption of cover crops. Comparing to fields in Heartland, those in Northern Crescent, Northern Great Plains, Prairie Gateway, and Mississippi Portal were less likely and those in Eastern Uplands and Southern Seaboard were more likely to adopt cover crops. Regional difference in adoption of cover crops or other conservation practices have been

found in other studies. Unger and Vigil (1998) find that the decision on adopting cover crops may vary from semiarid regions to humid/sub-humid regions depending on whether water is scarce or not. Some other studies (e.g., Davey and Furtan 2008, Ding, Schoengold, and Tadesse 2009, and Claassen et al. 2018) suggest that a higher rate of adoption in regions with greater rainfall may reflect the concern about soil erosion, while higher rate in dryer regions may reflect the importance of preserving soil moisture.

The likelihood of cover crop adoption was also affected by farmers' concerns. Farmers who had concerns over wind-driven erosion, soil compaction, water quality, or other concerns were more likely to adopt cover crops than those who did not have those concerns. The result aligns itself with the benefits that cover crops are supposed to provide. In contrast, farmers who had concerns over water-driven erosion, poor drainage, or low organic matter were less likely to adopt than those without such concerns. Steele, Coale, and Hill (2012) did not observe consistent differences in total organic matter and labile organic matter between the winter cover crop and control soils in an experiment with 13 years of cover crop use. Our finding on the effect of concerns over low organic matters is consistent with Steele, Coale, and Hill's (2012) finding. While plenty of studies have found positive effect of cover crops on soil organic matters and erosion by water (e.g., Shanks, Moore, and Sanders 1998; Ding et al. 2006; Dube, Chiduza, and Muchaonyerwa 2012;), many studies have concluded that cover crops' effect on organic matter may vary with cover crop species, soil type, and other practices, such as tillage and rotation (e.g., Wulanningtyas et al. 2021, Abdollahi and Munkholm 2014, Dube, Chiduza, and Muchaonyerwa 2012, and Motta et al. 2007). Our results imply that cover crops might either have not practically worked well on improving water-driven erosion, poor drainage and organic matter conditional on commonly used management and practices in soybean production or complexities of

management and implementation of cover crops as well as its interactions with other practices have discouraged its adoption to address those concerns.

Field characteristics also affected the adoption of cover crops. As expected, if a field was classified as "highly erodible", it was more likely to adopt cover crops; fields with slopes were also more likely to adopt cover crops since they with slopes are more vulnerable to soil erosion. Whether a field contained wetland did not statistically significantly affect the adoption. Soil texture also affected farmers' decision on cover crop adoption. Comparing to loam soil, fields with clay or mixed soil were more likely to adopt cover crops. There was, however, no statistically significant difference between loam soil and sandy or silty soil in cover crop adoption. Larger farms and larger fields were less likely to adopt cover crops. This is expected since the larger the farm/field, the more labor and time are needed for the implement and management of cover crops.

Farmer's demographic characteristics played an important role in practice adoption decision. Older farmers or farmers working at least 50% off farm were less likely to adopt cover crops. This might be due to the labor and time requirements of cover crop implementation and management as discussed previously. In addition, more educated farmers were more likely to adopt cover crops as they might better understand the importance of cover crops in environment and agricultural sustainability. Consistent with expectation, farmers who had land ownership were more likely to adopt cover corps as they care more about their own land. Drought in the last five years did not have an effect on the likelihood of cover crop adoption. This might be due to the same reason as what were discussed above. Cover crop's drought mitigating effect may interact with other factors such as cover crop species, planting and termination time, tillage, rotation, and soil type etc. Proper combinations of the use of cover crops and other practices

conditional on soil and weather conditions are required and many farmers might not have observed desired results on the use of cover crops as a drought adaptation strategy.

Results of Yield Moment Functions

Results of the first moment equations are presented in Table 4. As expected, more mGDD increased and more ODD decreased the soybean yield on average. If the year had drought in July or August, the mean of yield decreased. Regional difference showed in soybean yields. Fields in Heartland, which includes Illinois, Indiana, Iowa and parts of Missouri, Nebraska, South Dakota, Minnesota, Ohio, and Kentucky, had the highest yield among all regions. Not surprisingly, if soybeans were planted late or a bigger proportion of fields were replanted, the field had lower yield, resulting from shortened vegetative and reproductive intervals. Inputs of fertilizer did help increase yield. The more phosphorus was applied, the higher yield was. With small parameter for the quadratic term, the effect of phosphorus was close to linear. Application of manure also helped improve yield by adding more nutrients to the soil. Without cover crops, soybean yield in loam soil was higher than in clay, sandy, or mixed soil, but not significantly different from that in silty soil. This is consistent with the findings by previous studies such as Radocaj et al. (2020) and He et al. (2014).

Regional difference also showed in yield variance and skewness. Comparing to Heartland, soybean yield in Southern Seaboard had lower variations and that in Northern Great Plains, Prairie Gateway, Eastern Uplands, and Mississippi Portal had higher variations. In addition, Norther Crescent, Prairie Gateway, and Mississippi Portal had higher downside risk; in contrast, Northern Great Plains, Eastern Upland, and Southern Seaboard have lower downside risk. The higher percentage of field replanted, the more variation and the lower downside risk in yield, maybe caused by the replacement of damaged plants, for example by frosts. If soybeans

were planted late, their yield varied less but had higher downside risk, probably due to shorter growing season or higher probability of frost before harvest. At the mean application level, phosphorus inputs reduced the variation in yield, but manure application did not. In addition, more mGDD increased the variance, but more ODD decreased the variance. Both of them did not change the downside risk. As expected, drought occurred in July or August increased the variance of yield and in the meantime increased the risk of crop failure. Yield moments also showed heterogenous effect of soil texture.

Given the interaction terms with soil texture and weather, marginal effects of cover crops were calculated and presented in Table 5. Standard errors are calculated using delta method. As shown in Table 5, if there was no drought in July and August of the year, then cover crops statistically significantly increased the yield of soybeans planted in sandy, silty, or mixed soil; but there was no statistically significant effect on soybean yield in loam and clay soils. This is consistent with many studies and experiments that find cover crops help increase cash crop yield. In the meantime, cover crops increased yield variance in loam and clay soil, but decreased yield variance in sandy and silty soil and had no significant effect in mixed soil. In addition, planting cover crops reduced downside risk in loam soil while increased downside risk in all other types of soil. From above, we can see that effects of cover crops depend on soil type when there was no drought. And there was always a tradeoff among the mean, the variation, and the downside risk of yield, i.e., there was no simultaneous positive effect of cover crops on the three moments of yield.

With drought in July and August, cover crops reduced soybean yield in all soil types, although the effects in sandy and mixed soil were not statistically significant. This implies that cover crops consumed water in soil for their own growth and reduced water available for the

following cash crops. When drought occurred, the water supply was worsened to a point that crop yield decreased. In addition, cover crops increased yield variance but reduced the risk of crop failure in all types of soil when drought occurred. The mixture of positive and negative effect of cover crops on yield moments is somewhat consistent with the finding in the first stage that previous drought did not affect farmer's adoption of cover crops, implying a divided acceptance of cover crops as a drought adaptation strategy among farmers.

Conclusions

Through a 2SLS model we explored factors that affected farmer's adoption of cover crops by a logit model and examined the effects of cover crops on soybean yield and its risk by three moment functions. By incorporating two interplays between cover crops and soil type, and cover crops and drought, we were able to explore the varying effects of cover crops in drought and different soil types. While we found that the adoption of cover crops varied in regions and soil types and were affected by field properties and farmers' demographic characteristics and concerns, we did not find a significant effect of previous droughts on the adoption. This reveals a possibility that farmers were divided in the effect of cover crops in enhancing drought resilience. The results from the moment functions of soybean yield confirmed what the results in the first stage suggest. When there was a drought, cover crops reduced yield and increased yield variance. However, cover crops also reduced the downside risk or crop failure in the meantime. The mixed effects of cover crops on yield and its risk associated with an occurrence of drought support the statistically insignificant effect of previous drought on cover crop adoption, implying that farmers were divided in the acceptance of cover crops as a means to build resilience to drought. The mixed effect of cover crops also warrants a further study to calculate the certainty equivalent of net economic return of soybeans with cover crops, which requires information on cover crop

seed, planting, and termination cost as well as additional or saved fertilizer and pesticide cost. The certainty equivalent of net economic return of soybeans with cover crops may provide more information on economic impediments to farmer adoption of cover crops.

The low adoption rate of cover crops may also be related to complex interactions between management and cultural practices including species selection, planting and terminating date, rotation, and termination method. Achieving desired benefits can require significant training, learning and adjustments in many aspects of the farming system (Wallander et al. 2021). As shown in the National Cover Crop Survey 2020 (Conservation Technology Information Center 2020), roughly 70% of respondents said that they typically used their own experience or trial and error for cover cropping. About 67% and 60% of the respondents considered the two approaches, local farm tours to see how cover crops worked and one-on-one technical assistance to select, plant, or manage cover crops, very helpful or moderately helpful, respectively, in encouraging them to try cover cropping.

The recent increase in the cover crop adoption has been accompanied by financial incentives. In 2018, about one-third of the acreage planted with a cover crop received a financial assistance payment for cover crop adoption from either Federal, State, or other programs, ranging from $12 per acre to $92 per acre (Wallander et al. 2021). The USDA Natural Resources Conservation Service (NRCS) just announced a program to promote use of soil health practices especially cover crops. The initiative sets a goal of doubling the number of corn and soybean acres using cover crops to 30 million acres by 2030 (USDA 2022). Given the mixed effects of cover crops on soybean yield and yield risk as found in this study, financial incentives can help improve the certainty equivalent of net returns and thus encourage more risk-averse farmers to adopt cover crops.

We conclude the paper by recognizing a key limitation of this study is that it does not consider the long-term effects of sustainable practice adoption. For example, the payoffs from investments in improving soil fertility and reducing soil erosion are cumulative and long run and can reduce future expenses for crop nutrients, irrigation, and energy (Lee 2005; Tilman et al. 2001). By using cross-sectional data, this study does not capture such benefits; rather, a dynamic analysis using panel-data is needed to provide a more complete examination.

References

Abdollahi, L. and L. Munkholm 2014. Tillage system and cover crop effects on soil quality: I. Chemical, mechanical, and biological properties. *Soil Science Society of America Journal* 78(1): 262-270.

Acuña, J.C. and M. Villamil. 2014. Short-term effects of cover crops and compaction on soil properties and soybean production in Illinois. *Agronomy Journal* 106(3): 860-870.

Anderson, J. R., J. D. Dillion, and B. Hardaker. 1980. Agricultural Decision Analysis. Ames, Iowa: The Iowa State University Press.

Andraski, T.W. and L.G. Bundy. 2005. Cover crop effects on corn yield response to nitrogen on an irrigated sandy soil. *Agronomy Journal* 97(4): 1239-1244.

Antle, J. 1983. Testing the Stochastic Structure of Production: A Flexible Moment-Based Approach. *Journal of Business & Economic Statistics* 1(3): 192–201.

Antonakis, J., S. Bendahan, P. Jacquart, and R. Lalive. 2014. Causality and endogeneity: Problems and solutions. In D.V. Day (Ed.), The Oxford Handbook of Leadership and Organizations (pp. 93-117). New York: Oxford University Press.

Arbuckle, J. 2016. Iowa Farm and Rural Life Poll: 2015 Summary Report. https://dr.lib.iastate.edu/server/api/core/bitstreams/b8e999e7-f588-4dfb-9ed6-9ff93781f37a/content. Accessed on November 12, 2021.

Arora, V.K., C. Singh, A. Sidhu, and S. Thind. 2011. Irrigation, tillage and mulching effects on soybean yield and water productivity in relation to soil texture. *Agricultural Water Management* 98(4): 563-568.

Blanco-Canqui, H., T. Shaver, J. Lindquist, C. Shapiro, R. Elmore, C. Francis, and G. Hergert. 2015. Cover crops and ecosystem services: Insights from studies in temperate soils. *Agronomy journal* 107 (6): 2449-2474.

Butcher, K., A.F. Wick, T. DeSutter, A. Chatterjee, and J. Harmon. 2018. Corn and soybean yield response to salinity influenced by soil texture. *Agron. J.* 110: 1243-1253.

Claassen, R., M. Bowman, J. McFadden, D. Smith, and S. Wallander. 2018. Tillage intensity and conservation cropping in the United States. Economic Information Bulletin No. 197. Economics Research Services, USDA.

Clark, A., A. Decker, J. Meisinger, and M. McIntosh. 1997. Kill date of vetch, rye, and a vetch–rye mixture: I. Cover crop and corn nitrogen. *Agronomy Journal* 89(3): 427-434.

Conservation Technology Information Center (CTIC). 2015. Report of the 2014-2015 cover crop survey. Joint publication of the CTIC and the North Central Region Sustainable Agriculture Research and Education Program. West Lafayette, Indiana.

-2020. National Cover Crop Survey. https://www.ctic.org/files/CoverCropSurvey 2020 24mb.pdf. Accessed on January 15, 2022.

Corak, S. J., W. W. Frye, and M. S. Smith. 1991. Legume mulch and nitrogen fertilizer effects on soil water and corn production. *Soil Science Society of America Journal* 55 (5): 1395-1400.

Davey, K.A. and W. H. Furtan. 2008. Factors that affect the adoption decision of conservation tillage in the prairie region of Canada. *Canadian Journal of Agricultural Economics* 56(3): 257-275.

Dean, J.E. and R. Weil. 2009. Brassica cover crops for nitrogen retention in the Mid-Atlantic Coastal Plain. *Journal of environmental quality* 38(2): 520-528.

Day, R. H. 1965. Probability Distributions of Field Crops, *Journal of Farm Economics* 47: 713-741.

Di Falco, S. and J. Chavas. 2009. On Crop Biodiversity, Risk Exposure, and Food Security in the Highlands of Ethiopia. *American Journal of Agricultural Economics* 91(3): 599-611.

Di Falco, S., and M. Veronesi. 2014. Managing environmental risk in presence of climate change: the role of adaptation in the Nile Basin of Ethiopia. *Environmental and Resource Economics* 57(4): 553-577.

Ding, G., X. Liu, S. Herbert, J. Novak, D. Amarasiriwardena, and B. Xing. 2006. Effect of cover crop management on soil organic matter. *Geoderma* 130(3-4): 229-239.

Ding, Y. , K. Schoengold, T. Tadesse. 2009. The Impact of Weather Extremes on Agricultural Production Methods: Does drought increase adoption of conservation tillage practices? *Journal of Agricultural and Resource Economics* 34: 395-411.

Dong, F., D. Hennessy, and H. Jensen. 2010. Contract and exit decisions in finisher hog production. American Journal of Agricultural Economics 92(3): 667-684.

Dong, F., D. Hennessy, H. Jensen, and R. Volpe. 2016. "Technical efficiency, herd size, and exit intentions in US dairy farms." *Agricultural Economics* 47 (5): 533-545.

Dube, E., C. Chiduza, and P. Muchaonyerwa. 2012. Conservation agriculture effects on soil organic matter on a Haplic Cambisol after four years of maize–oat and maize–grazing vetch rotations in South Africa. *Soil and Tillage Research* 123: 21-28.

Ebelhar, S. A., W. W. Frye, and R. L. Blevins. 1984. Nitrogen from Legume Cover Crops for No-Tillage Corn 1. *Agronomy journal* 76(1): 51-55.

Groff, S., 2015. The past, present, and future of the cover crop industry. *Journal of Soil and Water Conservation* 70(6):130A-133A.

Hardaker, J.B., J. Richardson, G. Lien, and K. Schumann. 2004. Stochastic efficiency analysis with risk aversion bounds: a simplified approach. *Australian Journal of Agricultural and Resource Economics* 48(2): 253-270.

He, W., X. Luo, and G. Sun. 2014. The Trend of GIS-Based suitable planting areas for Chinese soybean under the future climate scenario. In Ecosystem Assessment and Fuzzy Systems Management; Cao, B.Y., Ma, S.Q., Cao, H.H., Eds.; Springer International Pubishing: Cham, Switzerland: 325–338.

Hunter, M., M. Schipanski, M. Burgess, J. LaChance, B. Bradley, M. Barbercheck, J. Kaye, and D. Mortensen. 2019. Cover Crop Mixture Effects on Maize, Soybean, and Wheat Yield in Rotation. *Agricultural and Environmental Letters* 4(1).

Jiang, P. and K. Thelen. 2004. Effect of Soil and Topographic Properties on Crop Yield in a North-Central Corn–Soybean Cropping System. *Agron. J.* 96: 252-258.

Karl, T. 1986. The Sensitivity of the Palmer Drought Severity Index and Palmer's Z-Index to their Calibration Coefficients Including Potential Evapotranspiration. *Journal of Climate and Applied Meteorology* 25 (1): 77–86.

Kaspar, T.C., D.J. Pulido, T. Fenton, T. Colvin, D. Karlen, D. Jaynes, and D. Meek. 2004. Relationship of Corn and Soybean Yield to Soil and Terrain Properties. *Agron. J.* 96: 700-709.

Kendall, M.G. and A. Stuart. 1977. The Advanced Theory of Statistics. Macmillan, New York, Vol. 1, 168.

Kessler, A., S. Archontoulis, and M. Licht. 2020. Soybean yield and crop stage response to planting date and cultivar maturity in Iowa, USA. *Agronomy Journal* 112:382–394.

Kim, K., Chavas, J.P., Barham, B., Foltz, J. 2014. Rice, irrigation and downside risk: a quantile analysis of risk exposure and mitigation on Korean farms. European Review of Agricultural Economics 41(5): 775–815.

Kravchenko, A.N., D. Bullock, and C. Boast. 2000. Joint Multifractal Analysis of Crop Yield and Terrain Slope. *Agron. J.* 92: 1279-1290.

Laloy, E. and C. Bielders. 2010. Effect of intercropping period management on runoff and erosion in a maize cropping system. *Journal of Environmental Quality* 39(3): 1001-1008.

Langdale, G. W., Blevins, R. L., Karlen, D. L., McCool, D. K., Nearing, M. A., Skidmore, E. L., Thomas, A. W., Tyler, D. D. and Williams, J. R. 1991. "Cover crop effects on soil erosion by wind and water". In Cover Crops for CleanWater. SWCS Edited by: Hargrove, W. L. 15–21. Ankeny, IA.

Lee, D. 2005. Agricultural Sustainability and Technology Adoption: Issues and Policies for Developing Countries. *American Journal of Agricultural Economics* 87: 1325-1334.

Linkemer, G., J. Board, and M. Musgrave. 1998. Waterlogging Effects on Growth and Yield Components in Late-Planted Soybean. *Crop Science* 38: 1576-1584.

Major, D. J., D. Johnson, J. Tanner, and I. Anderson. 1975. Effects of day length and temperature on soybean development. *Crop Science* 15: 174– 179.

McDaniel, M.D., L. Tiemann, and A. Grandy. 2014. Does agricultural crop diversity enhance soil microbial biomass and organic matter dynamics? A meta-analysis. *Ecological Applications* 24(3): 560-570.

Miguez, F.E. and G.A. Bollero. 2005. Review of corn yield response under winter cover cropping systems using meta-analytic methods. *Crop Science* 45(6): 2318-2329.

Miller, L., J. Chin, and K. Zook. 2012. Policy opportunities to increase cover crop adoption on North Carolina farms. Masters Project, Duke University, Durham, NC.

Mitchell J.P., A. Shrestha, and S. Irmak. 2015. Trade-offs between winter cover crop production and soil water depletion in the San Joaquin Valley, California. *Journal of Soil and Water Conservation* 70(6):430-440,

Motta, A. C., D. Wayne Reeves, C. Burmester, and Y. Feng. 2007. Conservation tillage, rotations, and cover crop affecting soil quality in the Tennessee valley: Particulate organic matter, organic matter, and microbial biomass. *Communications in soil science and plant analysis* 38(19-20): 2831-2847.

Munawar, A., R. Blevins, W. Frye, and M. Saul. 1990. Tillage and cover crop management for soil water conservation. *Agronomy Journal* 82(4): 773-777.

Muñoz, J.D., J. Steibel, S. Snapp, and A. Kravchenko. 2014. Cover crop effect on corn growth and yield as influenced by topography. *Agriculture, ecosystems & environment* 189:229-239.

National Drought Mitigation Center (NDMC). 2021. What Is USDM. https://droughtmonitor.unl.edu/About/WhatistheUSDM.aspx. Accessed on November 11, 2021.

National Research Council, 2007. Understanding American agriculture: challenges for the agricultural resource management survey. National Academies Press.

Nelson, K.A. and C. Meinhardt. 2011. Soybean Yield Response to Pyraclostrobin and Drainage Water Management. *Agronomy Journal* 103: 1359-1365.

Nielsen, D.C., D. Lyon, R. Higgins, G. Hergert, J. Holman, and M. Vigil. 2016. Cover crop effect on subsequent wheat yield in the central Great Plains. *Agronomy Journal* 108(1): 243-256.

OECD. 2011. Managing Risk in Agriculture: Policy Assessment and Design. Paris: Organization for Economic Cooperation and Development.

Paz, J., W. D. Batchelor, G. L. Tylka, and R. G. Hartzler. 2001. A Modeling Approach to Quantify the Effects of Spatial Soybean Yield Limiting Factors. *Transactions of the ASAE* 44(5): 1329–1334.

Pedersen, P., and M. Licht. 2014. Soybean growth and development. PM 19948. Ames, IA: Iowa State University Extension and Outreach.

Radočaj, D., M. Jurišić, V. Zebec, and I. Plaščak. 2020. Delineation of soil texture suitability zones for soybean cultivation: a case study in continental Croatia. *Agronomy* 10(6): 823.

Reddy, K. 2001. Effects of Cereal and Legume Cover Crop Residues on Weeds, Yield, and Net Return in Soybean (Glycine max). *Weed Technology* 15 (4): 660-668.

Reddy, P.P. 2017. Cover/Green Manure Cropping. In Agro-ecological Approaches to Pest Management for Sustainable Agriculture (pp. 91-107). Springer, Singapore.

Reuters. 2022. Off-season 'cover' crops expand as U.S. growers eye low-carbon future. https://www.reuters.com/markets/commodities/farming-climate-off-season-cover-crops-expand-us-growers-eye-low-carbon-future-2022-01-04/. Accessed on January 5, 2022.

Robertson, G. P., T. Bruulsema, R. Gehl, D. Kanter, D. Mauzerall, C. Rotz, and C. Williams. 2013. Nitrogen–climate interactions in US agriculture. *Biogeochemistry* 114(1): 41-70.

Sackett, J.L., 2013. An NCR-SARE cover crop project: Farmer-cooperator motivation and agronomic practices. *Journal of the NACAA* 6(2).

Schnepf, M. and C. Cox. 2006. Environmental benefits of conservation on cropland: the status of our knowledge. In Environmental benefits of conservation on cropland: the status of our knowledge, edited by M. Schnepf and C. Cox.

Shane, W. W., and K. Barker. 1986. Effects of Temperature, Plant Age, Soil Texture, and Meloidogyne incognita on Early Growth of Soybean. Journal of nematology 18(3): 320–326.

Shanks, L. W., D. Moore, and C. Sanders. 1998. Soil erosion. Cover cropping in vineyards. In A grower's handbook: 80-85.

Smith, R.G., L. Atwood, and N. Warren. 2014. Increased productivity of a cover crop mixture is not associated with enhanced agroecosystem services. *PloS one* 9(5): p.e97351.

Snapp, S.S., S. Swinton, R. Labarta, D. Mutch, J. Black, R. Leep, J. Nyiraneza, and K. O'neil. 2005. Evaluating cover crops for benefits, costs and performance within cropping system niches. *Agronomy journal* 97(1): 322-332.

Steele, M., F.J. Coale, R.L. Hill. 2012. Winter annual cover crop impacts on no-till soil physical properties and organic matter. *Soil Sci. Soc. Am. J.* 76:2164-2173.

Sustainable Agriculture Research and Education Program (SARE). 2017. Cover Crops. What is Sustainable Agriculture? UC Division of Agriculture and Natural Resources. https://sarep.ucdavis.edu/sustainable-ag/cover-crops. Accessed January 5, 2022.

Tack, J., A. Harri, and K. Coble. 2012. More than mean effects: Modeling the effect of climate on the higher order moments of crop yields. *American Journal of Agricultural Economics* 94(5): 1037-1054.

Tilman, D.K., J. Fargione, B. Wolff, C. D'Antonio, A. Dobson, R. Howarth, D. Schindler, W. Schlesinger, D. Simberloff, D. Swackhamer. 2001. Forecasting Agriculturally Driven Global Environmental Change. *Science* 292: 281-284.

Unger, P.W. and M. F. Vigil. 1998. Cover crop effects on soil water relationships. *Journal of soil and water conservation* 53(3):200-207.

USDA. 2022. USDA Offers Expanded Conservation Program Opportunities to Support Climate Smart Agriculture in 2022. https://www.usda.gov/media/press-releases/2022/01/10/usda-offers-expanded-conservation-program-opportunities-support. Accessed on January 15, 2022.

USDA Climate Hubs. 2021. Drought resistant practices. https://www.climatehubs.usda.gov/hubs/northeast/topic/drought-resistant-practices. Accessed on October 20, 2021.

USDA-ERS. 2000. Farm Resource Regions. Agricultural Information Bullentin Number 760. https://www.ers.usda.gov/webdocs/publications/42298/32489_aib-760_002.pdf?v=42487#:~:text=The%20Farm%20Resource%20Regions%20are%20derived%20from%20four,Land%20Resource%20Regions%2C%20and%20NASS%20Crop%20Reporting%20Districts. Accessed on September 5, 2021.

USDA-NRCS. 2018. Technical Note: Iowa Agronomy Technical Note 38 (Cover Crop Management). https://efotg.sc.egov.usda.gov/references/public/IA/Cover_Crop_Management_38_AGR_TN_2018_08.pdf. Accessed on November 1, 2021.

USDA Risk Management Agency. 2015. Conservation Compliance - Highly Erodible Land and Wetlands. Risk Management Agency Fact Sheet. https://www.rma.usda.gov/en/Fact-Sheets/National-Fact-Sheets/Conservation-Compliance-Highly-Erodible-Land-and-Wetlands#:~:text=Highly%20erodible%20land%20is%20any%20land%20that%20can,total%20field%20acreage%20that%20contains%20highly%20erodible%20soils. Accessed on January 15, 2022.

Vitale, J. D., C. Godsey, J. Edwards, and R. Taylor. 2011. The adoption of conservation tillage practices in Oklahoma: Findings from a producer survey. *Journal of Soil and Water Conservation* 66(4): 250-264.

Wallander, S., D. Smith, M. Bowman, and R. Claassen. 2021. *Cover Crop Trends, Programs, and Practices in the United States*. Economic Information Bulletin number 222, U.S. Department of Agriculture, Economic Research Service.

Wandel, J. and J. Smithers. 2000. Factors affecting the adoption of conservation tillage on clay soils in southwestern Ontario, Canada. American Journal of Alternative Agriculture 15(4): 181-188.

Westcott, P. and M. Jewison. 2013. Weather Effects on Expected Corn and Soybean Yields. FDS-13g-01. Economic Research Service/USDA.

Wulanningtyas, H., Y. Gong, P. Li, N. Sakagami, J. Nishiwaki, and M. Komatsuzaki. 2021. A cover crop and no-tillage system for enhancing soil health by increasing soil organic matter in soybean cultivation. *Soil and Tillage Research* Volume 205.

Table 1. Survey Means of Variables in the Logit Model

Variable and Unit	Mean	Std.Err
Cover crops (0/1)	0.100	0.003
Northern Crescent (0/1)	0.498	0.003
Northern Great Plains (0/1)	0.141	0.003
Prairie Gateway (0/1)	0.066	0.001
Eastern Uplands (0/1)	0.054	0.001
Southern Seaboard (0/1)	0.053	0.002
Mississippi Portal (0/1)	0.105	0.002
Northern Crescent (0/1)	0.083	0.001
Fields having moderate or steeper slope (0/1)	0.556	0.004
Soil texture: loam (0/1)	0.369	0.003
Soil texture: clay (0/1)	0.170	0.003
Soil texture: sandy (0/1)	0.069	0.002
Soil texture: mixed (0/1)	0.369	0.005
Soil texture: silty (0/1)	0.019	0.001
Concern about water-driven erosion (0/1)	0.262	0.003
Concern about wind-driven erosion (0/1)	0.080	0.002
Concern about soil compaction (0/1)	0.261	0.004
Concern about poor drainage (0/1)	0.232	0.004
Concern about low organic matter (0/1)	0.109	0.003
Concern about water quality (0/1)	0.066	0.002
Other concerns (0/1)	0.025	0.001
Field classified as "highly erodible" (0/1)	0.180	0.003
Field contains wetland (0/1)	0.041	0.002
Age (years)	57.730	0.100
Years of experience (years)	33.326	0.121
College education (0/1)	0.251	0.002
Off-farm work (0/1)	0.185	0.004
Land ownership (0/1)	0.491	0.004
Farm size (acres)	1348.206	11.558
drought5yr (years)	0.764	0.007
Field size (acres)	50.015	0.328

Table 2. Survey Means of Variables in Moment Functions

Variable and unit	Mean	Std. Err
Yield (bushels/acre)	52.354	0.097
Heartland (0/1)	0.498	0.003
Northern Crescent (0/1)	0.141	0.003
Northern Great Plains (0/1)	0.066	0.001
Prairie Gateway (0/1)	0.054	0.001
Eastern Uplands (0/1)	0.053	0.002
Southern Seaboard (0/1)	0.105	0.002
Mississippi Portal (0/1)	0.083	0.001
PlantLate (0/1)	0.129	0.002
ReplantPct	0.041	0.002
Manure (0/1)	0.043	0.001
Phosphorus (lbs/acre)	25.550	0.229
Labor (hours/acre)	1.503	0.017
drought18 (0/1)	0.175	0.002
Soil texture: loam (0/1)	0.369	0.003
Soil texture: clay (0/1)	0.170	0.003
Soil texture: sandy (0/1)	0.069	0.002
Soil texture: mixed (0/1)	0.369	0.005
Soil texture: silty (0/1)	0.019	0.001
Growing degree days (Celsius)	3304.675	4.396
Overheating degree days (days)	30.778	0.133
Cover crops	0.100	0.003

Table 3. Estimates of Parameters of the Logit Model

parameter	Estimation	Bootstrapped Std. Err.
Northern Crescent	-1.049***	0.157
Northern Great Plains	-3.100***	0.212
Prairie Gateway	-0.605***	0.205
Eastern Uplands	2.363***	0.129
Southern Seaboard	2.788***	0.172
Mississippi Portal	-1.187***	0.202
slope	0.520***	0.057
Soil texture: clay	0.711***	0.114
Soil texture: sandy	-0.249*	0.144
Soil texture: mixed	0.413***	0.087
Soil texture: silty	0.330	0.277
Concern about water-driven erosion	-0.413***	0.121
Concern about wind-driven erosion	0.532***	0.102
Concern about soil compaction	0.223*	0.129
Concern about poor drainage	-0.218**	0.105
Concern about low organic matter	-0.793***	0.121
Concern about water quality	0.901***	0.112
Other concerns	0.976***	0.142
Field "highly erodible"	0.545***	0.120
Field contains wetland	0.355	0.531
age	-0.024***	0.006
Years of experience	-0.005	0.005
College education	0.551***	0.078
Off-farm work	-0.258**	0.109
Land ownership	0.230*	0.080
Farm size	-9.280E-05***	3.160E-05
drought5yr	-0.089	0.068
Field size	-0.003***	0.001
Constant	-3.119***	0.667

Note: statistical significance at 1%, 5%, and 10% are denoted as ***, **, and *, respectively.

Table 4. Estimates of Parameters of Yield Moment Equations.

	First moment (Mean) equation		Second moment (variance) equation		Third moment (skewness) equation	
Parameter	Estimation	Bootstrapped Std. Err	Estimation	Bootstrapped Std. Err	Estimation	Bootstrapped Std. Err
Cover crops	-3.890	2.720	48.337***	9.092	2868.821***	341.341
Drought×cover crops	-20.160***	4.452	223.235***	21.604	16562.100***	1050.960
Cover crops×clay soil	6.869***	2.517	4.441	32.237	-12716.120***	1430.358
Cover crops×sandy soil	17.406***	2.509	-79.215***	17.073	-6299.486***	559.366
Cover crops×mixed soil	18.586***	1.655	-48.293***	18.088	-8944.720***	667.661
Cover crops×silty soil	14.110***	1.926	-154.608***	23.123	-4624.771***	793.497
Northern Crescent	-2.994***	0.314	-3.153	2.946	-263.873**	110.773
Northern Great Plains	-9.363***	0.472	19.329***	4.805	377.511***	146.329
Prairie Gateway	-4.436***	0.603	103.903***	7.312	-1081.599***	259.667
Eastern Uplands	-17.638***	1.147	39.311***	6.498	1982.181***	286.687
Southern Seaboard	-25.983***	0.819	-20.231***	4.326	932.423***	131.663
Mississippi Portal	-4.093***	0.668	153.902***	12.949	-2364.662***	500.023
PlantLate	-4.934***	0.256	-20.423***	-20.423***	-461.056***	77.673
ReplantPct	-6.639***	1.069	42.540***	11.625	4547.295***	499.821
Manure	1.956***	0.440	3.777	4.742	547.647***	184.636
Phosphorus	0.020***	0.004	-0.569***	0.062	-3.022	1.899
Phosphorus^2	1.735E-04***	2.680E-05	0.003***	0.001	0.003	0.012
Drought18	-3.696***	0.401	30.540***	4.323	-856.462***	151.164
Soil texture: clay	-1.507***	0.389	7.641**	3.275	-788.829***	113.448
Soil texture: sandy	-10.515***	0.477	71.824***	5.500	2274.254***	211.552
Soil texture: mixed	-3.742***	0.243	-17.207***	2.521	446.028***	84.746
Soil texture: silty	0.485	0.429	1.400	10.029	-1525.396***	389.240
mGDD	0.013***	0.001	0.041***	0.007	-0.314	0.242
ODD	-0.184***	0.016	-0.650***	0.189	-5.393	6.670
Constant	23.424***	1.390	7.472	18.158	1100.763*	648.716
R^2	0.334		0.095		0.112	

Note: statistical significance at 1%, 5%, and 10% are denoted as ***, **, and *, respectively.

Table 5. Marginal effects of cover crops under different situations

			yield equation		second moment (variance) equation		third moment (skewness) equation	
	Soil type	Drought	Estimation	Std.Err	Estimation	Std.Err	Estimation	Std.Err
cover crops	loam	No	-3.890	2.720	48.337***	9.092	2868.821***	341.341
		Yes	-24.049***	4.579	271.572***	26.449	19430.921***	1244.251
	Clay	No	2.980	3.108	52.778*	31.177	-9847.299***	1426.819
		Yes	-17.180***	3.826	276.013***	31.774	6714.801***	1166.486
	sandy	No	13.516***	2.349	-30.878**	13.859	-3430.665***	391.965
		Yes	-6.643	4.586	192.357***	23.436	13131.435***	1080.542
	mixed	No	14.696***	2.375	0.045	13.002	-6075.899***	461.852
		Yes	-5.463	4.216	223.280***	17.548	10486.201***	701.112
	silty	No	10.220***	1.839	-106.271***	21.328	-1755.950***	709.924
		Yes	-9.940***	4.234	116.964***	26.404	14806.150***	981.327

Note: statistical significance at 1%, 5%, and 10% are denoted as ***, **, and *, respectively.